



## LETTER TO EDITOR

# Pancytopenia: A Rare Presentation of Late Onset Isovaleric Acidemia

Hossein Karami<sup>1</sup>, Daniel Zamanfar<sup>2</sup>, Mohammad Reza Navaifar<sup>3</sup>, Parastoo Jamallivani<sup>4</sup>, Kimia Rasoli<sup>4</sup>, Mohammad Naderisorki<sup>1\*</sup>

<sup>1</sup>Thalassemia Research Center, Department of Pediatrics Hematology and Oncology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

<sup>2</sup>Diabetes Research Center, Department of Pediatrics, Bou Ali Sina Hospital, Mazandaran University of Medical Sciences, Sari, Iran

<sup>3</sup>Pediatric Infectious Diseases Research Center, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup>Student Research Committee, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

## ARTICLE INFO

### Article History:

Received: 27.11.2019

Accepted: 28.12.2019

### \*Corresponding author:

Mohammad Naderisorki,  
Bou Ali Sina Hospital, Mazandaran  
University of Medical Sciences,  
Pasdaran Boulevard, P.O. Box: 48158-  
38477, Sari, Iran  
Tel/Fax: +98 11 33342331  
Email: [dr.naderisorki@gmail.com](mailto:dr.naderisorki@gmail.com)

Please cite this article as: Karami H, Zamanfar D, Navaifar MR, Jamallivani P, Rasoli K, Naderisorki M. Pancytopenia: A Rare Presentation of Late Onset Isovaleric Acidemia. IJBC 2020; 12(1): 38-40.

## Dear Editor,

Isovaleric acidemia was first recognized by Tanaka and colleagues as a new genetic disorder of leucine metabolism in two siblings in early infancy. There is a defect in the catabolism of leucine resulting in the accumulation of isovaleric acid (1). Isovaleric acidemia (IVA) is an autosomal recessive inborn error of leucine metabolism caused by a deficiency of the mitochondrial enzyme isovaleryl-CoA dehydrogenase resulting in the accumulation of derivatives of isovaleryl-CoA. The clinical presentation of IVA is highly variable ranging from severely affected to asymptomatic individuals. It may present in the neonatal period as an acute episode of metabolic acidosis leading to coma and death or later as a chronic intermittent form associated with developmental delay and recurrent acidotic episodes during catabolic phases (2).

A three-year-old boy was admitted with vomiting, lethargy and loss of consciousness after a period of upper respiratory tract infection. The patient was the second child of non-relative parents. He was admitted nine months earlier with a similar presentation diagnosed as encephalitis. Upon admission, he was febrile and he had a pulse rate of 110/min and respiratory rate of 52/min. Physical examination was unremarkable for hepatosplenomegaly. Laboratory examination showed severe neutropenia, moderate anemia, and

thrombocytopenia and compensated metabolic acidosis.

Biochemical analysis including blood glucose and coagulation profile were within normal limits. Serum C-reactive protein, erythrocyte sedimentation rate and lactate dehydrogenase were normal. Serum ammonia level was 110 mcg/dL (19-90). Cerebrospinal fluid and urine analysis showed no abnormality. Blood and cerebrospinal fluid cultures were negative. The results of a complete blood count revealed the following: hemoglobin 8.7 g/dL; platelet count  $103 \times 10^9 / \mu\text{L}$ ; white blood cells  $0.8 \times 10^9 / \mu\text{L}$ ; neutrophil 70%; lymphocyte 17%; monocyte 5% and band cells 5%. Arterial blood gas showed pH 7.29;  $\text{Pco}_2$  27 mmHg;  $\text{Hco}_3$  13.4 mEq (22-26) with normal  $\text{Po}_2$  and  $\text{O}_2$  saturation. Urinary organic acid profile analyzed with Gas Chromatography Mass Spectrometry (GC-MS) revealed excessive excretion of isovalerylglycine (1320 times the normal value).

Due to persistent pancytopenia, bone marrow aspiration was performed which showed hypocellular marrow with trilineage hematopoiesis and relative monocytosis. Clinical findings of stupor and coma along with metabolic acidosis, mild hyperammonemia and increased excretion of urinary organic acids (isovalerylglycine) all were compatible with IVA. Upon diagnosis, low protein diet, L-carnitine and glycine were prescribed for the patient. After two weeks of treatment, metabolic acidosis resolved and white blood cells and platelets gradually normalized.

On day 14<sup>th</sup> of admission, laboratory data showed: WBC  $3.6 \times 10^9/\mu\text{L}$ , hemoglobin 12.8 g/dl; platelet count  $462 \times 10^9/\mu\text{L}$ ; pH 7.44; Pco<sub>2</sub> 37 mmHg and Hco<sub>3</sub> 25 mEq.

Severe pancytopenia has been previously reported in two infants with IVA. It was suggested that pancytopenia might be due to arrested maturation of hematopoietic precursors (3). There is also a report of IVA in a two-year-old female presenting as diabetic ketoacidosis who demonstrated neutropenia and thrombocytopenia.<sup>4</sup> Varying degrees of cytopenia in organic acidemia including IVA has been ascribed to the toxic effect of organic acids on hematopoietic cells. Secondary hemophagocytic syndrome has also been reported in IVA (3, 4). A newborn who was admitted at 10 days of age with lethargy, poor feeding, hypothermia, cholestasis and profound pancytopenia expired at 19 days of age. Autopsy showed extramedullary hematopoiesis and myelodysplasia of the bone marrow with arrest of the myeloid series at the promyelocytic stage. The appearance of bone marrow resembled promyelocytic leukemia; however, the 15:17 translocation was not present. The maturation arrest in granulopoiesis in IVA appears to be most likely due to a direct metabolic effect on granulocyte precursor cells (5).

Pancytopenia has been also reported in other types of organic acidemia such as methylmalonic and propionic acidemia. Bakshi and co-workers have reported three patients with MMA who presented with severe refractory pancytopenia during the acute illness. Their bone marrow examination revealed a wide spectrum of pathologies varying from bone marrow hypoplasia, hemophagocytosis to myelodysplasia with ring sideroblasts (6).

In a study on 46 children (4-5 years old) with various inherited metabolic disorders, different kinds of anemia were observed. Moreover, bicytopenia or pancytopenia was found in 8 (17.4%) of the children. The study indicated that in organic acidemias including methylmalonic acidemia, propionic acidemia, IVA, and argininosuccinic acidemia, most patients had anemia of chronic disease followed by vitamin B<sub>12</sub> deficiency and iron deficiency anemia. This study suggested that congenital anemias such as hereditary spherocytosis or thalassaemia could be observed as coexisting hematological diseases in young patients with inborn errors of metabolism (7).

An infant with propionic acidemia and reversible pancytopenia has been reported in whom light and electron bone marrow microscopy revealed severely disturbed cellular morphology with trilineage dysmyelopoiesis, hemophagocytosis and abundant multinucleated histiocytes. Evaluation of the infant's hematological abnormalities suggests that inhibition of bone marrow proliferation and maturation and shortened red blood cell survival were responsible for her pancytopenia. The studies performed in vitro suspect the role of propionic acid in this hematopoietic dysfunction (8). Severe neutropenia is also reported in an infant with methyl malonic acidemia (9). Also, three children with disorders of propionate metabolism have been presented, one with methylmalonic acidemia and two with propionic acidemia who developed secondary HLH during their

metabolic attacks (10).

In patients with IVA, hematological problems are frequently reported in early infancy and during acidotic episodes; hence cytopenia is not commonly encountered in later childhood (11). Our patient was a three-year-old boy whose diagnosis of IVA was confirmed at this age. Inborn errors of metabolism specifically late onset form of IVA should be considered in differential diagnosis of any patient with varying degrees of cytopenia and metabolic acidosis after the neonatal period.

**Patient Consent:** Written informed consent was obtained from the patients' parents for publication of this case report.

**Conflict of Interest:** None declared.

## References

1. Tanaka K, Budd M, Efron M, Isselbacher K. Isovaleric acidemia: a new genetic defect of leucine metabolism. *Proc Natl Acad Sci U S A*. 1966;56(1):236. doi: 10.1073/pnas.56.1.236. PubMed PMID: 5229850. PubMed Central PMCID: PMC285701.
2. Ensenaer R, Vockley J, Willard J-M, Huey JC, Sass JO, Edland SD, et al. A common mutation is associated with a mild, potentially asymptomatic phenotype in patients with isovaleric acidemia diagnosed by newborn screening. *The American Journal of Human Genetics*. 2004;75(6):1136-42. doi: 10.1086/426318. PubMed PMID: 15486829 PubMed Central PMCID: PMC1182150
3. Kelleher JF, Yudkoff M, Hutchinson R, August CS, Cohn RM. The pancytopenia of isovaleric acidemia. *Pediatrics*. 1980;65(5):1023-7. PubMed PMID: 7367115.
4. Kılıç M, Kaymaz N, Özgül RK. Isovaleric acidemia presenting as diabetic ketoacidosis: a case report. *J Clin Res Pediatr Endocrinol*. 2014;6(1):59. doi: 10.4274/Jcrpe.1181. PubMed PMID: 24637313. PubMed Central PMCID: PMC3986742.
5. Gilbert-Barness E, Barness LA. Isovaleric acidemia with promyelocytic myeloproliferative syndrome. *Pediatr Dev Pathol*. 1999;2(3):286-91. doi: 10.1007/s100249900125. PubMed PMID: 10191353
6. Bakshi NA, Al-Anzi T, Mohamed SY, Rahbeeni Z, AlSayed M, Al-Owain M, et al. Spectrum of bone marrow pathology and hematological abnormalities in methylmalonic acidemia. *American Journal of Medical Genetics Part A*. 2018;176(3):687-91. doi: 10.1002/ajmg.a.38599. PubMed PMID: 29330964
7. Tavit B, Sivri HSK, Coskun T, Gurgey A, Ozyurek E, Dursun A, et al. Haematological findings in children with inborn errors of metabolism. *Journal of Inherited Metabolic Disease: Official Journal of the Society for the Study of Inborn Errors of Metabolism*. 2006;29(5):607-11. doi: 10.1007/s10545-006-0379-8. PubMed PMID: 16906472
8. Stork LC, Ambruso DR, Wallner SF, Sambrano JE, Moscinski LC, Wilson HL, et al. Pancytopenia in propionic acidemia: hematologic evaluation and studies of hematopoiesis in vitro. *Pediatr Res*.

- 1986;20(8):783-8.doi: 10.1203/00006450-198608000-00017. PubMed PMID: 3737292
9. Guerra-Moreno J, Barrios N, Santiago-Borrero PJ. Severe neutropenia in an infant with methylmalonic acidemia. *Bol Asoc Med P R*. 2003;95(1):17-20. PubMed PMID: 12898746.
  10. Gokce M, Unal O, Hismi B, Gumruk F, Coskun T, Balta G, et al. Secondary hemophagocytosis in 3 patients with organic acidemia involving propionate metabolism. *Pediatr Hematol Oncol*. 2012;29(1):92-8.
  11. Malbora B, Avcı Z, Hasanoğlu A, Alehan F, Özbek N. Late onset of isovaleric acidemia presenting with bicytopenia. *Turk J Hematol*. 2010;27:216-8. doi: 10.5152/tjh.2010.34. PubMed PMID: 27263611