

Knowledge of Breast Cancer Risk Factors among a Cohort of Nurses in Iran

Ghahremanfard F¹, Ghorbani R^{2*}, Hemmati S³

1. Department of internal medicine, Semnan University of Medical Sciences, Semnan, Iran.

2. Research Center for Social Determinants of Health, department of community medicine, faculty of medicine, Semnan University of Medical Sciences, Semnan, Iran.

3. Semnan University of Medical Sciences, Semnan, Iran.

***Corresponding Author:** Ghorbani R, Email: ghorbani_raheb@yahoo.com

Submitted: 13-07-2014, Accepted: 08-08-2014

Abstract

Background: Improving the knowledge of healthcare providers towards different aspects of breast cancer leads to appropriate management and screening of disease. This study aimed to investigate the knowledge of breast cancer risk factors among a cohort of nurses in Iran.

Materials and Methods: The target group of the present study was 243 consecutive nurses (27 males and 216 females) at two referral hospitals in Semnan province, Iran, between March 2012 and February 2013. The data concerning the nurses' knowledge of breast cancer risk factors was collected using a predesigned questionnaire.

Results: In total, 9.9% of nurses had poor, 71.2% had intermediate and only 18.9% had good level of knowledge towards breast cancer risk factors. The level of knowledge towards risk factors was not affected by nurses' gender, age, level of education, years of experience, marital status, presence of a history of breast problem, or history of caring for a breast cancer patient. However, those nurses with a family history of breast cancer especially those with a patient as a first degree relative had significantly higher level of knowledge about the disease risk factors. The highest level of knowledge on breast cancer risk factors was observed among nurses working in CCU wards, while the lowest level of knowledge was found in pediatrics and psychiatry wards. Among nurses working in obstetrics and gynecology wards, 15.8% had a poor level, 68.4% had an intermediate level, and only 15.8% had an acceptable level of knowledge on breast cancer risk factors.

Conclusion: This study revealed a relatively low awareness of breast cancer risk factors among nurses emphasizing the need for adding proper education programs aimed at improving the knowledge of breast cancer risk factors among our nurses.

Keywords: Nurse, breast cancer, risk factor, knowledge.

Introduction

Breast cancer is the most common invasive malignancy among women and comprises about 23% of invasive cancers and 16% of all female cancers worldwide¹. The incidence of breast cancer varies greatly around the world with the lowest incidence in less-developed countries and the greatest incidence in the more-developed countries². However, because of poorer management and less appropriate controlling of risk factors, this cancer is a major public health issue in less developed countries leading to higher cancer-related death rates among women in these countries³. Moreover, although

the incidence of breast cancer has had a downward trend in developed countries within recent two decades, this trend has remained incremental in less developed areas which might be attributed to the transition toward a Western-style economy and improper pattern of reproductive behavior⁴. Also, increases in the proportion of women in the industrial workforce have affected the population distribution of established breast cancer risk factors⁵. Besides, some recent increased incidence of breast cancer in some high-risk Western countries may be due in part to greater use of

mammography and other specific and sensitive diagnostic tools^{6,7}. Iran as a developing country in Asia faces a high incidence rate of breast cancer and its-related mortality and morbidity, so that according to recent reports the incidence of breast cancer among women is 22 per 100,000 with the overall prevalence rate 120 per 100,000^{8,9}.

According to recent alarming statistics of breast cancer and its life-threatening consequences, improving the knowledge of healthcare providers towards different aspects of disease including what to look for, how to detect and diagnose breast cancer, and when the time is appropriate for its screening seems necessary¹⁰⁻¹². In this regard, great emphasis must be given to the issue of improving awareness of healthcare providers particularly nurses about breast cancer, its-related risk factors and screening methods. However, a few studies have been performed in Iran about nurses' knowledge towards different aspects of breast cancer. This study aimed to investigate the knowledge of breast cancer risk factors among a cohort of nurses in Iran.

Materials and Methods

In a cross-sectional study the target group which was 243 consecutive nurses (27 males and 216 females) working at two referral hospitals in Semnan province, Iran, between March 2012 and February 2013 were enrolled. The data was collected using a predesigned questionnaire. The questionnaire constituted of 19 questions about nurses' knowledge towards different aspects of breast cancer risk factors. The questionnaires were distributed and collected personally by the researcher after the respondents completed the filling process. The questionnaire consisted of two sections. The first section included personal information such as gender, age, education background, work experience, work specialization, working ward, history of providing health to breast cancer patients, and family history of breast cancer. The second section included information concerning the knowledge of nurses towards breast cancer risk factors. The respondents were requested to place a mark (X) on the choice they had selected. The level of nurses' knowledge was categorized as poor (the correct response to less than 50% of questions), intermediate (the correct response to 50% to 69.9% of questions), and good (the correct response

to more than 69.9% of questions). Results were reported as mean \pm standard deviation (SD) for the quantitative variables and percentages for the categorical variables. The groups were compared using the chi-square test (or Fisher's exact test if required) for the categorical variables. P values of 0.05 or less were considered statistically significant. All the statistical analyses were performed using SPSS version 21.0 (Armonk, NY: IBM Corp.).

Results

In total, 9.9% of nurses had poor, 71.2% had intermediate and only 18.9% had a good level of knowledge towards breast cancer risk factors. The overall level of knowledge towards this cancer was similar among male and female nurses, single and married nurses, younger and older ones, and also in those with different educational levels (Table 1). The level of knowledge was also not influenced by years of experience, presence of a history of breast problem, or a history of caring for a breast cancer patient. However, those nurses with a family history of breast cancer especially those with a patient as a first degree relative had significantly higher level of knowledge about disease risk factors. The highest level of knowledge on breast cancer risk factors was observed among nurses working in CCU wards followed by surgery ICU wards (Figure 1), while the lowest level of knowledge was found in pediatrics and psychiatry wards. Interestingly, among nurses working in obstetrics and gynecology wards, 15.8% had a poor level, 68.4% had an intermediate level, and only 15.8% had an acceptable level of knowledge of breast cancer risk factors.

Discussion

The main finding of the present study was an overall low level of knowledge among nurses towards breast cancer risk factors so that less than one-third of studied nurses had appropriate information about this subject and most of them had intermediate information about risk factors. More interestingly, in gynecology department with the most referrals of breast cancer patients, only 15.8% had an acceptable level of knowledge about breast cancer risk factors. Furthermore, intermediate to good level of knowledge was more observed in those nurses with a family history of breast cancer. These findings suggests that affected women with breast cancer who were referred

Table 1: Level of knowledge about breast cancer risk factors among participating nurses according to baseline characteristics

Characteristics	Level of knowledge		
	Poor	Intermediate	Good
Gender			
Men	5 (18.5)	19 (70.4)	3 (11.1)
Women	19 (8.8)	154 (71.3)	43 (19.9)
P-value	0.087		
Education level			
Non-college degree	2 (11.8)	11 (64.7)	4 (23.5)
College degree	22 (9.7)	162 (71.7)	42 (18.6)
P-value	0.827		
Age group			
< 30 y	12 (12.1)	71 (71.7)	16 (16.2)
30-39 y	7 (7.4)	69 (72.6)	19 (20.0)
≥ 40	5 (10.2)	33 (67.3)	11 (22.4)
P-value	0.294		
Marital status			
Single	5 (8.6)	43 (74.1)	10 (17.2)
Married	19 (10.3)	130 (70.3)	36 (19.5)
P-value	0.943		
Years of experience			
< 5 y	1 (12.9)	64 (68.8)	17 (18.3)
5-9 y	4 (5.6)	52 (73.2)	15 (21.1)
10-14 y	5 (14.3)	27 (77.1)	3 (8.6)
≥ 15 y	3 (6.8)	30 (68.2)	11 (25.0)
P-value	0.339		
History of breast problem			
Positive	1 (12.5)	5 (62.5)	2 (25.0)
Negative	23 (9.8)	168 (71.5)	44 (18.7)
P-value	0.852		
Family history of breast cancer			
Positive	0 (0.0)	20 (71.4)	8 (28.6)
Negative	24 (11.2)	153 (71.2)	38 (17.7)
P-value	0.038		
Care of a breast cancer patient			
Positive	9 (10.3)	59 (67.8)	19 (21.8)
Negative	15 (9.6)	114 (73.1)	27 (17.3)
P-value	0.592		




Figure 1: Level of knowledge about breast cancer risk factors according to different hospitals wards.

to these healthcare centers probably did not receive enough information or guidelines about their disease especially how to control and manage their risk factors. Similar results have been reported in some previous studies. In a study by Yaren et al., in Turkey, despite a high level of knowledge of breast cancer risk factors, inadequate knowledge of cancer screening methods was observed among nurses¹³. In another study by Ibrahim et al., in Nigeria, female doctors obtained a mean knowledge score of 74% and were the only professional group that had satisfactory knowledge of risk factors¹⁴. In a study by Chong et al., from Singapore, the majority of nurses had certain misconceptions in the knowledge of breast cancer and breast cancer screening. In this context, a higher proportion of those nurses in the family health services had higher knowledge scores than other nurses in the public health services¹⁵. In a study by Fotedar et al., in India, the average knowledge of breast cancer risk factors in the entire nursing population was only 49% so that 10.5% of nurses had poor knowledge, 25.2% of the nurses had good knowledge, 45% had very good knowledge and only 16.3% of the nurses had excellent knowledge about breast cancer risk factors and early detection methods¹⁶. Ahmed et al., in Pakistan found that only 37% of nurses had good knowledge of breast cancer risk factors¹⁷.

Thus, it seems that similar to other developing

countries discussed here, a relatively small proportion of our nursing population, even those who are working in healthcare centers providing caring services for breast cancer patients have a good knowledge of the breast cancer risk factors. One of the present study's limitations is that sometimes nurses had previously been employed in other wards or hospitals which could have a confounding effect.

Based on our results we think there is an urgent need to introduce more breast cancer education in nursing schools particularly in the public sector. Also results from the present study suggest the need for continuing medical education programs aimed at improving the knowledge of breast cancer risk factors among healthcare providers in our nursing population.

Conclusion

This study revealed a relatively low awareness of breast cancer risk factors among nurses emphasizing the need for adding proper education programs aimed at improving the knowledge of breast cancer risk factors among our nurses.

References

1. Byass P, de Courten M, Graham WJ, Laflamme L, McCaw-Binns A, Sankoh OA, et al. Reflections on the global burden of disease 2010 estimates. *PLoS Med.*

- 2013;10(7):e1001477.
2. Forouzanfar MH, Foreman KJ, Delossantos AM, Lozano R, Lopez AD, Murray CJ, et al. Breast and cervical cancer in 187 countries between 1980 and 2010: a systematic analysis. *Lancet*. 2011;378(9801):1461-84.
3. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, Thun MJ. Cancer statistics, 2008. *CA Cancer J Clin*. 2008;58(2):71-96.
4. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin*. 2011;61(2):69-90.
5. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin*. 2013;63(1):11-30.
6. Kelsey JL, Bernstein L. Epidemiology and prevention of breast cancer. *Annu Rev Public Health*. 1996;17:47-67.
7. Colditz GA, Rosner B. Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol*. 2000;152(10):950-64.
8. Mousavi SM, Montazeri A, Mohagheghi MA, Jarrahi AM, Harirchi I, Najafi M, et al. Breast cancer in Iran: an epidemiological review. *Breast J*. 2007;13(4):383-91.
9. Harirchi I, Ebrahimi M, Zamani N, Jarvandi S, Montazeri A. Breast cancer in Iran: a review of 903 case records. *Public Health*. 2000;114(2):143-5.
10. Sim HL, Seah M, Tan SM. Breast cancer knowledge and screening practices: a survey of 1,000 Asian women. *Singapore Med J*. 2009;50(2):132-8.
11. Seah M, Tan SM. Am I breast cancer smart? Assessing breast cancer knowledge among health professionals. *Singapore Med J*. 2007;48(2):158-62.
12. Lemlem SB, Sinishaw W, Hailu M, Abebe M, Aregay A. Assessment of Knowledge of Breast Cancer and Screening Methods among Nurses in University Hospitals in Addis Ababa, Ethiopia, 2011. *ISRN Oncol*. 2013;470981.
13. Yaren A, Ozkilinc G, Guler A, Oztop I. Awareness of breast and cervical cancer risk factors and screening behaviours among nurses in rural region of Turkey. *Eur J Cancer Care (Engl)*. 2008;17(3):278-84.
14. Ibrahim NA, Odusanya OO. Knowledge of risk factors, beliefs and practices of female healthcare professionals towards breast cancer in a tertiary institution in Lagos, Nigeria. *BMC Cancer*. 2009;9:76.
15. Chong PN, Krishnan M, Hong CY, Swah TS. Knowledge and practice of breast cancer screening amongst public health nurses in Singapore. *Singapore Med J*. 2002;43(10):509-16.
16. Fotedar V, Seam RK, Gupta MK, Gupta M, Vats S, Verma S. Knowledge of risk factors and early detection methods and practices towards breast cancer among nurses in Indira Gandhi Medical College, Shimla, Himachal Pradesh, India. *Asian Pac J Cancer Prev*. 2013;14(1):117-20.
17. Ahmed F, Mahmud S, Hatcher J, Khan SM. Breast cancer risk factor knowledge among nurses in teaching hospitals of Karachi, Pakistan: a cross-sectional study. *BMC Nurs*. 2006;5:6.