



LETTER TO EDITOR

Disseminated Intravascular Coagulation in a Case of Brucellosis Misdiagnosed as Thrombotic Thrombocytopenia Purpura

Nasim Valizadeh*

Assistant Professor of Hematology/Medical Oncology, Hematology-Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article History:

Received: 25.05.2015

Accepted: 30.08.2015

*Corresponding author:

Nasim Valizadeh, MD;

Assistant Professor of Hematology/

Medical Oncology, Hematology-

Oncology and Stem Cell

Transplantation Research Center,

Tehran University of Medical Sciences,

Tehran, Iran

Tel: +98 912 5474755

Fax: +98 21 88004140

Email: nsedaha0@gmail.com

Please cite this article as: Valizadeh N. Disseminated Intravascular Coagulation in a Case of Brucellosis Misdiagnosed as Thrombotic Thrombocytopenia Purpura. IJBC 2015; 7(5): 235-236.

Dear Editor

Brucellosis is a multisystem disease with various clinical presentations. It is a worldwide, zoonotic disease which has still remained as an uncontrolled health problem in many underdeveloped countries.¹ Hematological manifestations of brucellosis are so variable and include anemia, leukopenia, immune thrombocytopenia, thrombotic thrombocytopenia purpura (TTP) and hemophagocytic syndrome.²⁻⁷ Disseminated intravascular coagulation (DIC) has been reported in a few cases of brucellosis.⁸⁻¹⁰ Herein, we present a case of brucellosis who was presented with clinical picture of DIC misdiagnosed as TTP.

A 24-year-old woman was admitted with fever, abdominal pain and headache for the last three months. She was from a rural area in east Azarbaijan of Iran and had a history of fresh dairy consumption. On physical examination she had pallor, fever, and jaundice. Laboratory tests showed anemia (Hb: 7.6g/dl), thrombocytopenia (Platelet count $30 \times 10^9/\mu\text{l}$), indirect hyperbilirubinemia, and elevated lactate dehydrogenase (LDH). Peripheral blood smear revealed many fragmented RBCs. A diagnosis of TTP was made and urgent plasmapheresis was initiated for

the patient, but the results of further tests in the next few days showed prolonged prothrombin time (PT), increased fibrin degradation products (FDP) and D-dimer along with markedly elevated anti-brucella antibodies with Wright's and Coombs Wright's tests, thus the diagnosis of DIC secondary to brucellosis was affirmed. Treatment with doxycycline and rifampin was initiated for the patient. Three days following appropriate antimicrobial therapy thrombocytopenia and coagulopathy were resolved and she was discharged with anti-brucellosis medication.

Although TTP is characteristically defined by a pentad of thrombocytopenia, microangiopathic hemolytic anemia, fluctuating neurological signs, renal impairment and fever, it can be considered without fulfilling the whole pentad. In another word positivity for only the first 3 above mentioned items could be present in approximately 75% of the cases.⁵ TTP is a thrombotic microangiopathy similar to DIC; however, in contrast to DIC, the mechanism of thrombosis in TTP is not via the coagulation pathway activation and consumptive coagulopathy. The results of the blood coagulation assays (PT, PTT, D-Dimer, FDP and fibrinogen levels) in TTP were normal in our patient.

According to the presence of fever, headache,

microangiopathic hemolytic anemia and thrombocytopenia, diagnosis of TTP was made for our patient, but further laboratory studies indicating prolonged PT, increased FDP and D-dimer was compatible with the diagnosis of DIC secondary to brucellosis.

There are various hematological manifestations in brucellosis which include anemia, leucopenia, thrombocytopenia, TTP, hemophagocytic syndrome and rarely DIC.²⁻⁹ As a result, in patients with brucellosis presenting with abnormal hematological features such as DIC, diagnosis of the disease may be problematic and delayed, accordingly we should consider brucellosis in any patient with unexplained DIC in endemic areas for brucellosis.

Conflict of Interest: None declared.

References

1. Mantur BG, Amarnath SK. Brucellosis in India- a review. *J Biosci*. 2008;33:539-47. PMID: 19208979.
2. Citak EC, Citak FE, Tanyeri B, Arman D. Hematologic manifestations of brucellosis in children: 5 years experience of an anatolian center. *J Pediatr Hematol Oncol*. 2010;32(2):137-40. doi: 10.1097/MPH.0b013e3181ced382.
3. Yilmaz M, Tiryaki O, Namiduru M, Okan V, Oguz A, Buyukhatipoglu H, et al. Brucellosis-induced immune thrombocytopenia mimicking ITP: a report of seven cases. *Int J Lab Hematol*. 2007;29(6):442-5. PMID: 17988299.
4. Sevinc A, Buyukberber N, Camci C, Buyukberber S, Karsligil T. Thrombocytopenia in brucellosis: case report and literature review. *J Natl Med Assoc*. 2005;97(2):290-3. PMID: 15712797.
5. Kiki I, Gundogdu M, Albayrak B, Bilgiç Y. Thrombotic thrombocytopenic purpura associated with *Brucella* infection. *Am J Med Sci*. 2008;335(3):230-2. doi: 10.1097/MAJ.0b013e3180d09f19.
6. Di Mario A, Sica S, Zini G, Salutari P, Leone G. Microangiopathic hemolytic anemia and severe thrombocytopenia in *Brucella* infection. *Ann Hematol*. 1995;70(1):59-60. PMID: 7827209.
7. Valizadeh N, Musavi J, Nikoonejad AR, Nateghi Sh. Acute prostatitis and hemophagocytic syndrome in a case of Brucellosis. *Shiraz E-Med J*. 2011;12(2):107-11.
8. Turunc T, Demiroglu YZ, Kizilkilic E, Aliskan H, Boga C, Arslan H. A case of disseminated intravascular coagulation caused by *Brucella* melitensis. *J Thromb Thrombolysis*. 2008;26(1):71-3. PMID: 17562127.
9. Akbayram S, Dogan M, Akgun C, Peker E, Parlak M, Oner AF. Disseminated intravascular coagulation in a case of brucellosis. *Clin Appl Thromb Hemost*. 2011;17(6):E10-2. doi: 10.1177/1076029610378501.
10. Almér LO. A case of brucellosis complicated by endocarditis and disseminated intravascular coagulation. *Acta Med Scand*. 1985;217(1):139-40. PMID: 3976429.