

# Iranian Journal of Blood & Cancer

Journal Home Page: www.ijbc.ir

#### **ORIGINAL ARTICLE**

# Cognitive, Emotional, and Behavioral Problems of Children with Hemophilia

# Manijeh Firoozi\*

Department of Clinical Psychology, University of Tehran (Pardis Pharabi), Ghom, Iran

#### ARTICLE INFO ABSTRACT Article History: Background: Children with hemophilia are prone to a variety of psychological Received: 16.01.2017 problems due to some limitations associated with the disease. We aimed to Accepted: 10.04.2017 compare the cognitive, emotional, and behavioral problems of children with hemophilia to healthy children. Kevwords: Methods: This study was performed on 65 children with hemophilia and 65 Hemophilia healthy individuals as the control group who were between the ages of 7 and Behavioral disruption 12 years in Children's Hospital. The Child Behavior Checklist (CBCL) was used Emotional problem to identify emotional/behavioral problems and Wisconsin Card Sorting Test Executive function (WCST) to evaluate cognitive problems. Child behavior checklist Results: The results showed that children with hemophilia obtained lower scores Wisconsin card sorting test in activity, academic performance, and overall competence variables. Children with hemophilia in comparison to healthy children showed more internalizing \*Corresponding author: and externalizing problems and emotional and behavioral deficits. Also they Manijeh Firoozi, demonstrated more impairment in executive functions than healthy children. No.401, Department of Clinical Conclusion: The bio-psycho-social factors such as factors associated with the Psychology, University of Tehran disease (e.g. anemia and bleeding), and the treatment (e.g. side effects of the (Pardis Pharabi), Ghom, Iran Tel: +98 25 36166100 drugs) and environmental and social factors are among underlying causes of Email: mfiroozy@ut.ac.ir some psychological problems in children with hemophilia. Please cite this article as: Firoozi M. Cognitive, Emotional, and Behavioral Problems of Children with Hemophilia. IJBC 2017; 9(3): 69-74.

#### Introduction

Hemophilia is an inherited bleeding disorder linked to the X chromosome resulting in a genetic deficiency in clotting factor VIII (hemophilia A) or factor IX (hemophilia B). It is classified as mild, moderate, or severe considering the percentage of the factor that is produced.<sup>1</sup> Patients with hemophilia are at increased risk of joint bleeding (typically in the knees, elbows and ankles), intra peritoneal and intracranial hemorrhage, bleeding during and after surgery. There are also complications due to natural course of the chronic disease and the treatments the patients have to receive such as pain, anxiety, and depression.<sup>2</sup> Children with hemophilia encompass a difficult life since they have to deal with chronic arthropathy followed by fatigue and limb limitations. One of the serious concerns of parents and health care systems is emotional and behavioral problems in children with hemophilia which gets more complicated with increasing age and may affect their quality of life.<sup>3</sup>

Chronic diseases can lead to emotional, behavioral, and cognitive problems.<sup>4</sup> Children with chronic diseases confront extreme stress, deprivation and serious limitations associated with their disease.<sup>5</sup> Perceived stress can interfere with brain functions and may result in emotional and behavioral dysfunction. Hanson and colleagues indicated that difficult life circumstances in childhood can decrease hippocampus and amygdala volume which cause behavioral disorders.<sup>6</sup> Children with hemophilia are often deprived of regular life activities because of the fear of recurring episodes of bleeding.<sup>7</sup> The protracted stress can make them vulnerable to the brain atrophy which may result in behavioral problems.<sup>8</sup> Prolonged bleeding causes a feeling of fatigue and weakness in children with hemophilia. The vitality of children will be reduced and it will provide a context for developing a sense of anger and frustration.<sup>9</sup> Children with hemophilia are susceptible to various infectious diseases and viruses such as hepatitis and AIDS.<sup>10</sup> Parents of these children show overprotective behavior towards their children, resulting in anxiety and chronic depression in the children.<sup>11</sup> Numerous studies have emphasized on low compliance of young patients with hemophilia to medical instructions. The emotional and behavioral problems disrupt the compliance with the treatment.<sup>12</sup>

Psychological consequences of hemophilia are worth studying. In this study, we aimed to assess the psychological status of children with hemophilia in three behavioral, emotional and cognitive aspects. Assessment of behavioral disorders and cognitive impairment in children with hemophilia by Wisconsin Card Sorting Test (WCST) is being investigated for the first time in our study comparing with healthy children.

#### **Materials and Methods**

The present research was a descriptive causalcomparative study. The study population consisted of all boys with hemophilia (aged 7-12 years) who referred to Children's Hospital. Healthy children were selected as the control group from siblings of the patients. The convenience sampling method was applied for the study. The exclusion criteria were as follows: 1) existence of comorbidity, 2) history of psychiatric disorders, 3) having the ability to participate in psychological interventions, and 4) presence of any kind of intracranial hemorrhage. The effect of gender variable was also controlled due to differences in incidence of behavioral and emotional disorders between girls and boys.

The sample consisted of 65 healthy children and 65 children with hemophilia who were assessed through Child Behavior Checklist (CBCL) and WCST. Informed consent was taken from the parents in advance explaining the motives of the study.

#### Child Behavior Check List (CBCL)

The Child Behavior Checklist (CBCL) is a report provided by the caregiver about children and is widely used in research and clinical practice. Generally, CBCL consists of the several following parts: 1) Academic performance assessment of children and adolescents in the fields of cognitive ability, training, and education issues, 2) Social skills assessment of children and adolescents for evaluating their adaptation with peers, siblings, parents and how they cope with challenges, and 3) Assessment of emotional and behavioral problems in children and adolescents.

The internal consistency of the CBCL was estimated using Cronbach's alpha coefficients. Alpha coefficient of competence was relatively high and its range was between 65- 85% for CBCL. The test reliability based on Achenbach experimental assessment approach and by using Cronbach's alpha was 0.89 for boys and 0.94 for girls and by using split-half reliability was 0.84 for boys and 0.87 for girls. The results of the construct validity supported the 8-factor structure of this scale by using factor analysis in Iran. In addition, convergent validity of this scale with "Junior Eysenck Personality Questionnaire (JEPQ)" and "Rutter behavioral problems questionnaire" was satisfactory.

If the index of discrimination power which is related to the variance and distribution scores reach higher than 90%, the scale will be appropriate. Obtained coefficients in the CBCL form were all at the high level. Based on the mentioned information about validity, it can be concluded that, CBCL is a valid tool in assessing behavioral and emotional problems and can be used with confidence by users.<sup>13</sup>

#### Wisconsin Card Sorting Test (WCST)

The Wisconsin Card Sorting Test (WCST) which was first developed by Berg and Grant is a useful tool for studying cognitive deficits after brain injuries.<sup>14</sup> WCST has been widely used as a neuropsychological test of "setshifting", i.e. the ability to display cognitive flexibility and abstract reasoning. Participants must maintain a concept found at the stage of testing in sequential conditions. When the classification rules change, they must change the previous concepts. Subjects were given a set of 64 cards which there were 1 to 4 symbols on them as triangles, stars and circles in 4 colors: red, green, yellow, and blue. Of course, there were no two identical cards. Four cards including a "red triangle", two "green stars", three "yellow Plus sign" and four "blue circles" were used as the main cards. The participant was told to match the cards based on 4 main patterns on the cards. After each response, participants received right or wrong feedback. In fact, whether a particular match is right or wrong. Scores obtained from this test included: number of incorrect responses, perseverative errors, and percentiles of achieved categories. Validity of this test has been reported 0.86 for measurement of cognitive deficits after brain injury.

The Kolmogorov-Smirnov test was applied to test normal distribution. It indicated normal distribution of Externalizing/Conduct, Inattention/Hyperactivity, Internalizing/Emotional and Social/Peer variables in experience-based scales. There was no normal distribution in other variables. Therefore, independent T-test was used to analyze the findings of mentioned variables and other variables were analyzed by The Mann-Whitney test.

#### Results

In this study, children with hemophilia (n=65) were compared to healthy children (n=65) in terms of cognitive, emotional, and behavioral problems (Table 1). The analysis of covariance demonstrated that there were no significant differences in age and socioeconomic status between the two groups. The minimum and maximum age in children with hemophilia was 7.5 to11.2 years and in control group it was was 7.8 to 11.6 years. Most of the children in both groups were in third and fourth grade of elementary school. Most parents were educated at the university level. The severity of hemophilia in most patients were of moderate type.

As shown in Table 2, there was no significant difference

Table 1: Demograp	hic data of children with hemophi	lia and healthy children		
Variable		Children with Hemophilia	Healthy children (siblings)	%
		Μ	Μ	
Age of the child (m	in, max)	7.5,11.2	7.8,11.6	
School grade				
	1 <sup>th</sup>			0.5
	2 <sup>th</sup>			2.1
	3 <sup>th</sup>			43.6
	4 <sup>th</sup>			41.2
	5 <sup>th</sup>			12.6
Mother's	Diploma and under			23
education	University graduated			77
Father's education	Diploma and under			35
	University graduated			65
Type of hemophilia	L			
	Mild		_	12.7
	Moderate		_	69.4
	Sever			18.6

Table 2: Mann-Whitney test to compare the behavioral-emotional problems in CBCL

Variables	Children	Children with Hemophilia		Healthy children (siblings)		P value
	Μ	SD	Μ	SD	Whitney U	
Social functions	6.11	1.72	6.54	2.21	671	0.43
Academic Achievement	1.31	4.68	5.55	0.79	455	0.002
Anxiety/Depression	4.4	6.67	3.05	2.96	361	0.001
Withdrawal/Depression	2.67	3.33	1.57	1.18	369	0.001
Somatic complaints	2.05	2.17	1.3	0.92	472	0.004
Thought problems	3.83	4.23	1.91	1.38	355	0.001
Attention deficit	4.65	6.1	3.16	3.53	464	0.003
Aggressive behavior	8.93	11.47	4.72	5.03	372	0.001
Internalization	7.61	12.17	5	4.88	321	0.001
Affective disorder	3.77	4.87	1.62	1.79	321	0.001
Anxiety disorder	2.45	3.41	1.54	1.7	398	0.001
Somatization	1.27	0.61	0.46	0.88	722	0.721
ADHD	4.15	5.26	3.12	2.97	524	0.024
Oppositional defiant disorder (ODD)	2.31	2.43	0.98	1.22	482	0.005

between the two groups in average score of social behavior (P=0.430). The average score of academic performance in children with hemophilia was significantly lower than the healthy control group (P<0.01). Average scores of children with hemophilia in anxiety/depression, withdrawal/ depression, somatic complaints, social problems, thought problems, attention problems, rule-breaking behavior, aggressive behavior, internalizing problems, emotional problems, anxiety, attention deficit hyperactivity disorder, and oppositional behavior were significantly higher than healthy children, while there was no significant difference between the two groups in physical problems (P=0.727).

According to Table 3, the result of t-test indicated that the average scores of activities and general competence variables in hemophilic children were significantly lower than control group and externalization and behavioralemotional problems were significantly higher in children with hemophilia than healthy children (P>0.01).

As shown in Table 4, the average scores of the percentiles of achieved categories in children with hemophilia was significantly lower and preservation error variable was higher than healthy children.

According to Table 4, t-test results indicated that the average score of incorrect responses in children with hemophilia were significantly higher than healthy children (P>0.01).

#### Discussion

Based on the findings in our study, children with hemophilia indicated significantly lower scores than the control group in externalization/conduct, inattention/ hyperactivity, internalization/emotions, and academic performance. This results were in line to the study of Coppola and colleagues who showed the academic achievement in children and adolescents with hemophilia were significantly lower than healthy children.<sup>15</sup> Children with hemophilia would be more absent from the school due to occasional bleeding episodes. On the other hand, physical limitations prevent them from participating in school activities and do their homework same as their healthy peers. This could explain their poor academic achievement.

Table 3: T-test to compare activitie	s, general competence,	externalizing and general	problems according to CBCL form

Problem Scales	Children with Hemophilia		Healthy children (siblings)		t	P value	
	Μ	SD	Μ	SD			
Externalizing/Conduct	16.24	5.13	6.28	2.18	4.93	0.001	
Inattention/Hyperactivity	12.79	6.8	8.53	2.07	3.16	0.002	
Internalizing/Emotional	17.6	4.09	11.62	2.72	3.34	0.001	
Social/Peer	11.62	5.3	9.2	3.01	2.69	0.06	
Total	49.73	13.18	20.93	7.22	5.13	0.001	

Table 4. T-test	of Wisconsin	Card Sorting	Test (	WCST	) for	nartici	nants
1able 4. 1-lest	. OI WISCONSIII	Caru Sorting	5 1051 (	WCDI	101	particip	Jams

Problem Scales	Children with H	emophilia	Healthy children (siblings)		t	P value
	Μ	SD	Μ	SD	-	
Categories Achieved	4.16	0.93	5.4	1.65	3.76	0.012
Failure to Maintain Set	3.67	2.5	5.17	4.06	4.15	0.001
Trial to first Category	2.36	2.91	5.32	3,45	4.97	0.001
Trial administered	15.8	4.31	19.2	2.43	2.43	0.03
Total trial Correct	16.1	3.55	20.43	2.64	2.67	0.01

The results of our study showed that social behaviors in children with hemophilia do not differ from healthy children. Chiu and colleagues investigated the social functioning of children with hemophilia; they showed there were no significant differences in popularity and social acceptance of children with hemophilia compared to their healthy classmates.<sup>16</sup> Chronic diseases cause increase in social sensitivity in the patients. They will also be concerned about judgment of their peers, so they comply with normal social behavior of their community to avoid negative self-image. As a result, they suppress their feelings and accept norms more easily.

Obviously, children with hemophilia compared with healthy children show more anxious/depressed and withdrawn/depressed conditions, somatic complaints, thought problems, attention problems and aggressive behavior, internalizing and externalizing problems and generally behavioral-emotional problems. Khair and colleagues conducted a qualitative study on 30 children with hemophilia.<sup>17</sup> They revealed that hemophilia had a significant effect on family lives, educational issues, school and traveling plans. Most of them felt frustrated and expressed anger against the disease. Tryzapch and colleagues also assessed emotional functioning in hemophiliac children.<sup>18</sup> As their reports, children with hemophilia showed more problems in internalizing and anxiety/depression disorders. Children with hemophilia showed symptoms of depression more commonly compared to healthy children.19 There was also higher level of anxiety in hemophiliac children which was significantly correlated with the parents' attitude.<sup>20</sup> Children with hemophilia in comparison to healthy group displayed more affection issues, attention deficit hyperactivity disorder and oppositional defiant disorder. A four-year longitudinal study indicated that children with hemophilia experienced more emotional problems; however, their quality of life improved over the time.<sup>21</sup>

Current research findings indicate a cognitive impairment in children with hemophilia through Wisconsin Card Sorting Test (WCST). WCST as a neuropsychological test displays frontal lobe functioning and cognitive flexibility (for assessment of executive functions, behavioral regulation, and social discourse). In a systematic review, Janual and colleagues demonstrated that hemophilia could not cause per se cognitive deficit and reduced IQ, but adverse effects of the disease could lead to serious cognitive and behavioral damages.<sup>22</sup> So, we consider the bio-psycho-social factors as the main reason for behavioral and emotional problems.

Experience of pain and anemia effect on mood and temper of children with hemophilia (as bio factor). Over-parenting, lack of social learning and negative self-perception are involved in emotional problems (as a psychological factor). Children with hemophilia are less prone to socialization because they are afraid of participating in group activities. They are at increased risk for rejection by peers. So, social isolation may be a trigger for depression and anxiety (as a social factor). On the other hand, the child may feel anger followed by environmental and social deprivations. It would be expressed as aggressive behavior and conduct disorder.

In the study of Tryzapch and colleagues, children with hemophilia had not more externalizing problems compared to healthy children.<sup>23</sup> However, in this study, externalizing problems in hemophiliac children were more observed than healthy children. This difference may be explained by cultural and social issues, parenting styles and parents' exaggerated understanding of child behavior problems.

The results of our study showed children with hemophilia demonstrate more impairment in executive functions. Although, previous studies have shown children with hemophilia who had a history of intracranial hemorrhage experienced impaired intelligence and visual perception, there was no cases of intracranial hemorrhage in our study. We showed cognitive function is influenced by the disease itself even in the absence of an intracranial hemorrhage.

Restriction in physical and social activities of hemophiliac children from early childhood may interfere with development of cognitive processes. Also, many cognitive skills do not shape in these children properly. Frequent absence from the school may be another factor that makes children to be less involved in school activities and it provides an inappropriate context for stimulation and training of intelligence and cognitive skills. This research suggests to perform future investigation of various psychological and neurological aspects in children with hemophilia in larger populations of the patients.

# Conclusion

The bio-psycho-social factors such as factors associated with the disease (e.g. anemia and bleeding), and the treatment (e.g. side effects of the drugs) and environmental and social factors are among underlying causes of some psychological problems in children with hemophilia.

# Conflict of Interest: None declared.

### References

- Nathwani AC, Tuddenham EG, Rangarajan S, Rosales C, McIntosh J, Linch DC, et al. Adenovirus-associated virus vector-mediated gene transfer in hemophilia. N Engl J Med. 2011; 22(365):2357-65. doi: 10.1056/ NEJMoa1108046. PubMed PMID: 22149959. PubMed Central PMCID: PMC3265081.
- Nathwani AC, Reiss UM, Tuddenham EG, Rosales C, Chowdary P, McIntosh J, et al. Long-term safety and efficacy of factor IX gene therapy in hemophilia B. N Engl J Med. 2014;371(21):1994-2004. doi: 10.1056/NEJMoa1407309. PubMed PMID: 25409372. PubMed Central PMCID: PMC4278802.
- Srivastava A, Brewer AK, Mauser-Bunschoten EP, Key NS, Kitchen S, Llinas A, et al. Guidelines for the management of hemophilia. Haemophilia. 2013; 19(1):e1-47. doi: 10.1111/j.1365-2516.2012.02909.x. PubMed PMID: 22776238.
- Miller GE, Chen E, Parker KJ. Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms. Psychol Bull. 2011; 137(6):959-97. doi: 10.1037/a0024768. PubMed PMID: 21787044. PubMed Central PMCID: PMC3202072.
- Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and agerelated disease. Physiol Behav. 2012; 106(1):29-39. doi: 10.1016/j.physbeh. PubMed PMID: 21888923.
- Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. Nat Rev Neurosci. 2012; 13(10):701-12. doi: 10.1038/nrn3346. PubMed PMID: 22968153.
- Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and agerelated disease. Physiol Behav. 2012; 106(1):29-39. doi: 10.1016/j.physbeh.2011.08.019. PubMed PMID: 21888923.
- Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and agerelated disease. Physiol Behav. 2012; 106(1):29-39. doi: 10.1016/j.physbeh.2011.08.019. PubMed PMID: 21888923.
- 9. Duggal P, Thio CL, Wojcik GL, Goedert JJ, Mangia

A, Latanich R, et al. Genome-wide association study of spontaneous resolution of hepatitis C virus infection: data from multiple cohorts. Ann Intern Med. 2013; 158(4):235-45.doi:10.7326/0003-4819-158-4-201302190-00003. PubMed PMID: 23420232.

- James SR, Nelson K, Ashwill J. Nursing care of children: Principles and practice. 4th ed. Saunders; 2014
- Coppola A, Tagliaferri A, Di Capua M, Franchini M. Prophylaxis in children with hemophilia: evidencebased achievements, old and new challenges. Semin Thromb Hemost. 2012; 38(1):79-94. doi: 10.1055/s-0031-1300954. PubMed PMID: 22314606.
- Achenbach TM, Ruffle TM. The child behavior checklist and related forms for assessing behavioral/ emotional problems and competencies. Pediatr Rev. 2000; 21(8):265-71. PubMed PMID: 10922023.
- Monchi O, Petrides M, Petre V, Worsley K, Dagher A. Wisconsin Card Sorting revisited: distinct neural circuits participating in different stages of the task identified by event-related functional magnetic resonance imaging. J Neurosci. 2001; 21(19):7733-41. PubMed PMID: 11567063.
- Coppola A, Tagliaferri A, Di Capua M, Franchini M. Prophylaxis in children with hemophilia: evidencebased achievements, old and new challenges. Semin Thromb Hemost. 2012; 38(1):79-94. doi: 10.1055/s-0031-1300954. PubMed PMID: 22314606.
- Coppola A, Tagliaferri A, Di Capua M, Franchini M. Prophylaxis in children with hemophilia: evidencebased achievements, old and new challenges. Semin Thromb Hemost. 2012; 38(1):79-94. doi: 10.1055/s-0031-1300954. PubMed PMID: 22314606.
- Noll RB, Gartstein MA, Vannatta K, Correll J, Bukowski WM, Davies WH. Social, emotional, and behavioral functioning of children with cancer. Pediatrics. 1999; 103(1):71-8. PubMed PMID: 9917442.
- Khair K, Meerabeau L, Gibson F. Self-management and skills acquisition in boys with hemophilia. Health Expect. 2015; 18(5):1105-13. doi: 10.1111/hex.12083. PubMed PMID: 23711015.
- Franchini M, Mannucci PM. Past, present and future of hemophilia: a narrative review. Orphanet J Rare Dis. 2012; 7:24. doi: 10.1186/1750-1172-7-24. PubMed PMID: 22551339. PubMed Central PMCID: PMC3502605.
- Valentino LA, Mamonov V, Hellmann A, Quon DV, Chybicka A, Schroth P, et al. A randomized comparison of two prophylaxis regimens and a paired comparison of on-demand and prophylaxis treatments in hemophilia A management. J Thromb Haemost. 2012; 10(3):359-67. doi: 10.1111/j.1538-7836.2011.04611.x. PubMed PMID: 22212248. PubMed Central PMCID: PMC3488301.
- Berntorp E, Shapiro AD. Modern haemophilia care. Lancet. 2012; 379(9824):1447-56. doi: 10.1016/S0140-6736(11)61139-2. PubMed PMID: 22456059.
- 21. Broderick CR, Herbert RD, Latimer J, Barnes C, Curtin JA, Mathieu E, et al. Association between

physical activity and risk of bleeding in children with hemophilia. JAMA. 2012; 308(14):1452-9. doi: 10.1001/jama.2012.12727. PubMed PMID: 23047359.

22. Tchanturia K, Davies H, Roberts M, Harrison A, Nakazato M, Schmidt U, et al. Poor cognitive flexibility in eating disorders: examining the evidence using the Wisconsin Card Sorting Task. PLoS One. 2012; 7(1):e28331. doi: 10.1371/journal.

pone.0028331. PubMed PMID: 22253689. PubMed Central PMCID: 3257222.

 Revel-Vilk S, Golomb MR, Achonu C, Stain AM, Armstrong D, Barnes MA, et al. Effect of intracranial bleeds on the health and quality of life of boys with hemophilia. J Pediatr. 2004; 144(4):490-5. doi: 10.1016/j.jpeds.2003.12.016. PubMed PMID:15069398.