Zinc Status and its Correlation with Basic Parameters in Transfusion Dependent Thalassemic Patients: A Pakistani Perspective

Sultan S *1, Irfan SM 1, Kaker JU 1, Zeeshan R 1, Kidwai A 1

- 1. Liaquat National Hospital, Karachi, Pakistan.
- 2. Afzal Memorial Thalassemia Foundation, Karachi, Pakistan.

*Corresponding Author: Sultan S, Email: sadia.sultan@Inh.edu.pk

Submitted: 18-12-2013, Accepted: 20-02-2014

Abstract

Background: β-thalassemia is an autosomal hemoglobinopathy with inconsistent universal distribution. Among patients with thalassemia diverse non-siderotic complications distinctly influence the attribute of life, including zinc deficiency due to varied etiologies.

The objective of the present study was to determine zinc levels in patients with β-thalassemia major and its correlation with maternal characteristics, hematological parameters, liver enzymes, serum ferritin, duration of chelation and number of transfusions among Pakistani patients.

Patients and Methods: Seventy-five β-thalassemia major patients on desferrioxamine were enrolled from August 2010 to July 2012. CBC, liver function tests, serum ferritin, HbsAg and Anti-HCV were evaluated. Zinc level was measured using atomic absorption spectrophotometer.

Results: The mean age of patients was 10.57 ± 3.5 years. Forty one (54.7%) and 34(45.3%) patients were males and females respectively. The frequency of zinc deficiency was 24%. We established positive correlation between zinc deficiency and longer duration of chelation (P<0.001) and also with anemia (P<0.001). No correlation could be established with other parameters.

Conclusion: Our study revealed that hypozincemia is not unusual in β -thalassemic patients on desferrioxamine. We propose zinc levels should be regularly measured primarily in anemic patients with long duration of chelation.

Keywords: β-thalassemia major, zinc deficiency, desferrioxamine.

Introduction

β-thalassemia major symbolizes a group of recessively inherited hemoglobinopathies, first depicted by Cooley and Lee and exemplified by diminished synthesis of β globin chains 1 . The homozygous state typically needs medical care within the first year of life and is manifested by severe anemia that necessitates unremitting transfusion for continued survival 1 , 2 . Appropriate combination of blood transfusion support and iron chelation remedies has noticeably lengthened the life expectancy among these patients to the fourth decade of life 1 , 3 .

Nevertheless iron overload is still a major culprit of mortality and morbidity among thalassemics³. Cardiac impairment secondary to iron accumulation and oxidative injury represent the major cause of

mortality, which can be averted by chelation ³. In the last four decades, the conventional drug used for iron chelation has been desferrioxamine (DFO). DFO has obviously improved the prospects of this disease as it is a powerful iron chelator, prevents iron accumulation and iron mediated organ damage ⁴. But on the other hand it also chelates certain important minerals such as zinc, cobalt and copper ⁴.

Zinc is a crucial trace mineral needed by all living tissues because of its decisive requirement in proteins synthesis and also acting as a cofactor in enzyme catalysis ^{5, 6}. Zinc is found in nearly all cells and has fundamental role in immune system involving both innate and acquired immunity ⁶. Zinc is also actively involved in many metabolic

Sultan et al.

activities in human body such as DNA synthesis, cellular growth, wound healing, fertility and conception ^{5, 6}. Beside it also has antioxidant properties thus protecting the cells from free radicals injuries, which are frequent finding in thalassemics, and is also involved in host defense mechanism such as chemotaxis of granulocytes ⁶. The significance of zinc in human body is illustrated by the sound effects of zinc deficiency which includes impaired wound healing, osteoporosis, delayed skeletal maturation, growth retardation, brittle hairs, impaired glucose metabolism, impaired sense of smell and taste, reduced fertility, neurological disturbance, irritability and deformed nails ^{3-5, 7}.

Zinc deficiency in thalassemia is ascertained owing to various etiologies which integrate oral or parenteral iron chelation, inadequate diet, and augmented urinary zinc excretion and ongoing hemolysis ^{1, 4, 8, 9}.

In thalassemia major trace metal zinc deficiency has been debatable 3. Many regional and international studies give conflicting results. Studies reported from Iran, Iraq and Egypt have revealed pronounced zinc deficiency in contrast to another study reported from Tehran, Iran which has revealed high levels of zinc in patients on DFO ^{1, 2, 8, 10}. Another study from Kuwait has reported zinc levels in thalassemia major patients to be within normal limits ¹¹.

Some studies recommend performing routine zinc assessment in thalassemic patients and to replenish it when required ⁴. Currently no consensus for regular monitoring and no meticulous guidelines or recommendations for screening of these patients exists. However, it is well proven that zinc supplementation has a beneficial role and has been shown to enhance linear growth among thalassemic patients ⁷.

The primary objective of our study was to evaluate the serum zinc levels in Pakistani transfusion dependent thalassemic patients on DFO. Whilst secondary objective was to ascertain any correlation of zinc levels with maternal characteristics, hematological parameters, biochemical markers, duration of chelation and number of transfusions.

Patients and Methods:

Patients

This was a prospective study extending from August 2010 to July 2012. Seventy five, non consecutive

patients with established β -thalassemia major were enrolled in this study. Thirty patients were in regular follow up prior to study; others were registered during the study period. Ethical endorsement was given by ethical and research committee of Liaquat National Hospital. An informed consent was attained from parents or Patients' guardians since all patients were under 18 years old.

All registered patients were on regular blood transfusions and parenteral iron chelation. Patients on oral iron chelation, having chronic liver disease (child score 2 and 3), diarrhea, malabsorption and those suffering from acute infections were excluded. Patients with β thalassemia intermedia and compound heterozygous form were also excluded from the study.

Methods

Demographic data including age and gender (maternal characteristics), number of blood transfusions and duration of chelation were recorded by taking thorough history from parents or patients' guardians and from patients' medical records. Blood samples for accessing the zinc levels were collected prior to blood transfusion under aseptic conditions. To perform this 5cc of peripheral venous blood in sodium heparin bottle was taken from the patient and zinc levels were assessed using an atomic absorption spectrophotometry. The defined reference value for zinc was 50-150 μ g/dl, zinc deficiency was defined zinc serum level of <50 μ g/dl.

Hematological parameters including hemoglobin, hematocrit, WBC and platelets were determined by automated Sysmex analyzer XT 2000i (Sysmex, Japan). Serum ferritin and liver function tests were performed by HITACHI 912 using a photometric assay. Hepatitis B and hepatitis C infections were detected by chemiluminescence method (Abbott Axsym, USA).

Statistical Analysis

Data was entered and analyzed using SPSS software (Version 17, SPSS Inc., Chicago, Illinois, USA). The results were expressed as mean ± SD for quantitative variables and qualitative variables were presented as frequency and percentages. Student's t-test was applied for the comparison of means. Data were considered statistically significant at P value < 0.05. Correlation analysis was performed using

the spearman correlation method to identify relationship between the serum zinc levels and maternal characteristics, duration of chelation, number of transfusions, hematological markers and serum ferritin. Chi- square test was applied for evaluating the correlation between zinc levels and hepatitis B and C status.

Results:

In this prospective study, 75 patients with β -thalassemia major were enrolled. Among these patients 41 (54.7%) were males and 34 (45.3%) were females. The mean age was 10.57 \pm 3.5 (range 5 to 15) years. The mean serum zinc level was 71.45 \pm 21.98 (range 30.4-117.0). The ratio of insufficiency was higher in male gender by a 2:1 ratio (male: female).

Overall 18 patients 24% were zinc deficient, out of which 12 were male and 6 were female. The patients were divided into two groups: zinc deficient and non zinc deficient. The mean zinc levels in zinc deficient and non deficient groups were 44.03 ± 5.48 and 80.11 ± 17.64 respectively (P < 0.001). Baseline characteristics of both zinc deficient and non deficient patients are shown in table 1.

Zinc deficiency also revealed positive correlation with low hemoglobin (0.566, 0.014) as shown in figure 1. The mean hemoglobin was 8.94±1.11 in non deficient group, whereas was significantly lower, 6.01±0.67 in the deficient group (Figure 2) (P<0.001). Zinc deficiency also had positive

correlation with higher duration of chelation being 7.72±3.06 (years) and 2.95± 1.45 (years) in zinc deficient and non deficient patients respectively (P<0.001).

Serum ferritin were significantly higher in all patients but no significant correlation was detected with zinc deficiency (P>0.05). HbsAg was found to be positive in 2.7% while Anti-HCV was reactive in 32% of patients. There were no statistically significant correlations (P>0.05) between low levels of zinc and maternal characteristic, TLC and platelets count, elevated liver enzymes, hepatitis B, hepatitis C and number of transfusions per month.

Discussion:

 β -thalassemia major causes decreased erythrocytes survival and iron burden owing to repeated blood transfusions and diverse pathological expressions of primary and secondary complications. The key treatment strategy is to decrease the iron overload which enhances the life expectancy of these patients.

The current study investigated the status of serum zinc in thalassemic patients on parenteral iron chelation. The explanation for zinc deficiency is the zinc binding affinity of desferrioxamine which augments urinary zinc elimination, hyperzincuria culminating in gradual zinc depletion in thalassemics ^{4, 12-14}.

Zinc is a key constituent which is essential for maintenance of cellular homeostasis. The principal

Table 1: Demographic findings among zinc deficient and non deficient thalassemic patients.

Evaluated parameters	Deficient group Serum zinc < 50 μg / dl Mean±SD	Non deficient group Serum zinc ≥ 50 Mean±SD	<i>P</i> value
Age	11.56±3.51	10.26±3.46	0.1
Serum zinc μg / dl	44.03± 5.48	80.11± 17.64	< 0.001
Duration of chelation(years)	7.72± 3.06	2.95± 1.45	< 0.001
Number of transfusion(per month)	2.22±0.42	2.19±0.97	0.9
Hemoglobin gm/dl	6.01±0.67	8.94±1.11	< 0.001
Hematocrit(%)	19.42±3.76	26.57±3.46	< 0.001
Ferritin µg / dl	4058.89±2340.02	4040.5±2090.07	0.9
Total bilirubin mg/dl	1.35±1.12	1.43±1.13	0.7
Direct bilirubin	0.55±0.66	0.58±0.67	0.8
Indirect bilirubin	0.78±0.55	0.82±0.59	0.7
AST u/l	55.83±38.63	59.26±34.69	0.7
ALT u/I	54.44±45.72	67.91±64.18	0.4




Figure 1: Positive correlation of zinc deficiency with low hemoglobin levels (P<0.001).

reason for its mandatory requirement is the fact that hundred of metallo enzymes require zinc as a cofactor to be functionally effective. Zinc insufficiency is deemed as one of the core causative factors for impaired linear growth, delayed sexual maturity, neuropsychological changes, immune dysfunction and endocrinopathies in thalassemics individuals ^{3,5,7,15}.

Diverse studies have revealed considerably diminished zinc levels in thalassemia major patients. Al Samarrai et al. from Iraq demonstrated zinc deficiency among thalassemic patients and attributed it to hyperzincuria 16. Shamshirsaz et al. have reported 79.6% of their thalassemic patients to have zinc deficiency and subsequent delayed sexual activity, retarded growth and significant association with lumbar BMD 1. The authors emphasized that the cause of zinc deficiency was insufficient zinc intake which could be attributed to a high prevalence of deficiency of this trace mineral in general Iranian population 1. Analogous results have been reported from Egypt, by Nasr et al. in their series of 64 patients². They reported that the mean zinc level was significantly lower (12.4±5.4µg/dl) in patients versus the control group (95.1±10.3µg/dl). They also analyzed association of zinc insufficiency with duration of transfusion and chelation therapy

but no correlation were established, with sub optimal chelation ². In contrast our results shows statistically significant correlation (P<0.001) of zinc deficiency with duration of chelation and signifies the enhanced clearance associated with increased duration of chelation. Augmented urinary zinc excretion in patients receiving desferrioxamine supports our finding ¹⁴.

The present study illustrated that hypozincemia is not uncommon in thalassemics regardless of age. Similarly a report by Dehshal et al. showed that 37% of thalassemic patients had zinc deficiency and authors proposed that serum zinc levels be routinely monitored in these patients 4. Likewise Ferdaus et al. from Bangladesh also found that desferrioaxime treatment was associated with zinc loss in 60% of their thalassemic patients³. Analogous to above mentioned studies, another regional study, by Ghone et al. reported from India showed diminished levels of serum zinc in their series of 72 patients ¹⁷. A study on Iranian thalassemic patients have reported 10% of thalassemic patients to have zinc insufficiency while 52% of those had variable degree of depression 18,19. Another study has identified a reverse correlation of zinc deficiency with blood transfusion rate (P < 0.05) 18. Also Bekheirnia et al. from Iran found 84.8% of




Figure 2: Comparison of hemoglobin levels in zinc deficient and non deficient groups.

thalassemic patients to be zinc deficient¹⁹. Our findings are also in concurrence with studies reported from Turkey, Tunisia, Thailand and Egypt ²⁰⁻²³.

In contrast to our results Bashir from Jordan found significantly increased zinc levels 24 in their thalassemic patients. In parallel Mehdizadeh et al. from Tehran also reported significantly higher serum zinc levels in 64 thalassemic patients ¹⁰. The regularity of blood transfusion appears to somehow, prevent such deficiency as a consequence of receiving healthy blood donation and it may clarify the normal or even elevated serum zinc levels ^{10,12}. As in our series, zinc deficiency was detected in under transfused patients with low hematocrit. Al Awadhi et al. found normal levels of serum zinc in 49 thalassemia major patients which correlates in their study with non compliance of patients with chelation therapy 11. Similarly Kwan and colleagues reported only 3 out of 68 thalassemic patients to have zinc deficiency ²⁵. Kajanachumpol et al. reported plasma and hair zinc levels to be much lower but erythrocytes' zinc was higher in thalassemic patients than controls. The mechanism of increased erythrocytic zinc was not clear but may be the impact of impaired zinc metabolism ²². Zinc deficiency in thalassemics has been debatable and controversial. The disparities in different reports can be accredited to regional factors, variable dietary habits, various chelation agents,

non compliance to chelation, difference in age and also transfusional practices.

Conclusion:

Our study revealed that hypozincemia is not unusual in β -thalassemic patients on desferrioxamine. We propose zinc levels should be regularly measured primarily in anemic patients with long duration of chelation.

Acknowledgments:

We sincerely thank the Department of Statistics, Liaquat National Hospital for biostatistical analysis of this manuscript. The authors are grateful to all parents and patients participating in this study. We are also thankful to clinical staff and laboratory services for their collaboration in the study.

References:

- Shamshirsaz AA, Bekheirnia MR, Kamgar M, Pourzahedgilani N, Bouzari N, Habibzadeh M, et Al. Metabolic and endocrinologic complications in beta-thalassemia major: a multicenter study in Tehran. BMC Endocr Disord. 2003;3(1):4.
- Nasr MR, Ali S, Shaker M, Elgabry E.. Antioxidant micronutrients in children with thalassaemia in Egypt. East Mediterr Health J. 2002; 8(4-5):490-5.
- 3. Ferdaus MZ, Hasan AK, Shekhar HU. Analysis of serum lipid profiles, metal ions and thyroid hormones levels abnormalities in beta-thalassaemic

Sultan et al.

- children of Bangladesh. J Pak Med Assoc. 2010; 60(5):360-4.
- Dehshal MH, Hooghooghi AH, Kebryaeezadeh A, Kheirabadi M, Kazemi S, Nasseh A, et al. Zinc deficiency aggravates abnormal glucose metabolism in thalassemia major patients. Med Sci Monit. 2007; 13(5): 235-9.
- Ackland ML, Michalczyk A. Zinc deficiency and its inherited disorders -a review. Genes Nutr. 2006; 1(1):41-9.
- Shazia Q, Mohammad ZH, Rahman T, Shekhar HU. Correlation of oxidative stress with serum trace element levels and antioxidant enzyme status in Beta thalassemia major patients: a review of the literature. Anemia. 2012;2012:270923.
- Arcasoy A, Cavdar A, Cin S, Erten J, Babacan E, Gözdasoglu S, et al. Effects of zinc supplementation on linear growth in beta-thalassemia (a new approach). Am J Hematol. 1987;24(2):127-36.1987; 24(2):127-36.
- 8- Erdoğan E, Canatan D, Ormeci AR, Vural H, Aylak F. The effects of chelators on zinc levels in patients with thalassemia major. J Trace Elem Med Biol. 2013; 27(2):109-11.
- Fikry SI, Saleh SA, Sarkis NN, Mangoud H. Study of serum zinc in relation to nutritional status among thalassemia patients in Damanhour. J Egypt Public Health Assoc. 2003;78(1-2):73-93.
- 10. Mehdizadeh M, Zamani G, Tabatabaee S. Zinc status in patients with major beta-thalassemia. Pediatr Hematol Oncol. 2008; 25(1):49-54.
- Al-Awadhi AM, Alfadhli SM, Al-Khaldi D, Borhama M, Borusly M.. Investigation of the distribution of lymphocyte subsets and zinc levels in multitransfused beta-thalassemia major patients. Int J Lab Hematol. 2010; 32(2):191-6.
- 12. Rashidi M, Aboomardani M, Rafraf M, Arefhosseini SR, Keshtkar A, Joshaghani H.. Effects of Vitamin E and Zinc Supplementation on Antioxidants in Beta thalassemia major Patients. Iran J Pediatr. 2011; 21(1):8-14.
- Uysal Z, Akar N, Kemahli S, Dincer N, Arcasoy A..
 Desferrioxamine and urinary zinc excretion in beta-thalassemia major. Pediatr Hematol Oncol. . 1993; 10(3):257-60.
- 14. Aydinok Y, Coker C, Kavakli K, Polat A, Nisli G, Cetiner N, et al. Urinary zinc excretion and zinc status of patients with beta-thalassemia major. Biol Trace Elem Res. 1999;70(2):165-72.
- 15. Consolini R, Calleri A, Legitimo A, Massei F.

- Immunological evaluation of patients with beta-thalassemia major. Acta Haematol.. 2001;105(1):7-12.
- Al-Samarrai AH, Adaay MH, Al-Tikriti KA, Al-Anzy MM. Evaluation of some essential element levels in thalassemia major patients in Mosul district, Iraq. Saudi Med J. 2008; 29(1):94-7.
- 17. Ghone RA1, Kumbar KM, Suryakar AN, Katkam RV, Joshi NG. Oxidative stress and disturbance in antioxidant balance in beta thalassemia major. Indian J Clin Biochem. 2008;23(4):337-40.
- 18. Moafi A1, Mobaraki G, Taheri SS, Heidarzadeh A, Shahabi I, Majidi F. Zinc in thalassemic patients and its relation with depression. Biol Trace Elem Res. 2008;123(1-3):8-13.
- Bekheirnia MR1, Shamshirsaz AA, Kamgar M, Bouzari N, Erfanzadeh G, Pourzahedgilani N, et al. Serum zinc and its relation to bone mineral density in beta-thalassemic adolescents. Biol Trace Elem Res.2004; 97(3):215-24.
- Kassab-Chekir A1, Laradi S, Ferchichi S, Haj Khelil A, Feki M, Amri F, et al. Oxidant, antioxidant status and metabolic data in patients with beta-thalassemia. Clin Chim Acta. 2003; 338(1-2):79-86.
- Arcasoy A, Canata D, Sinav B, Kutlay L, Oguz N, Sen M. Serum zinc levels and zinc binding capacity in thalassemia. J Trace Elem Med Biol. 2001; 15(2-3):85-7.
- 22. Kajanachumpol S1, Tatu T, Sasanakul W, Chuansumrit A, Hathirat P.. Zinc and copper status of thalassemic children. Southeast Asian J Trop Med Public Health. 1997;28(4):877-80.
- 23. Fuchs GJ1, Tienboon P, Linpisarn S, Nimsakul S, Leelapat P, Tovanabutra S Nutritional factors and thalassaemia major. Arch Dis Child. 1996;74(3):224-7.
- 24. Bashir NA. Serum zinc and copper levels in sickle cell anaemia and beta-thalassaemia in North Jordan. Ann Trop Paediatr. 1995;15(4):291-3.
- 25. Kwan EY1, Lee AC, Li AM, Tam SC, Chan CF, Lau YL, et al. A cross-sectional study of growth, puberty and endocrine function in patients with thalassaemia major in Hong Kong. J Paediatr Child Health. 1995;31(2):83-7.