Preparation of Plasminogen by Affinity Chromatography

Mousavi Hosseini K^{1*}, Nikougoftar Zarif M¹

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran.

*Corresponding Author: Mousavi Hosseini K, Email: mkmousavi@yahoo.com

Submitted: 12-04-2014 , Accepted: 15-05-2014

Abstract

Background: Plasminogen is one of the compounds derived from human plasma. Activation of plasminogen produces plasmin. Plasmin is able to lyse fibrinogen, fibrin, and some other human plasma proteins. The aim of the present work was to study the separation of human plasminogen by affinity chromatography using gel lysine Sepharose.

Materials and Methods: Normal human plasma was used as the starting material. By increasing its concentration of ethanol from 0% to 8% in pH 7.2 at -3°C the centrifugation was carried out and the supernatant I was separated from precipitate I. Then supernatant fraction I was used for separation of human plasminogen from plasma and further purification was carried out using affinity chromatography using affinity gel lysine Sepharose.

Results: In our study we could obtain plasminogen with the concentration of 0.5%. In normal human plasma the concentration of plasminogen is about 200mg/l. Comparing the result with the concentration of plasminogen in normal human plasma shows about 25 folds higher concentration.

Conclusion: Using affinity chromatography method a 25-fold increase in plasminogen concentration was achieved which shows this method is a promising method for the preparation of plasminogen from human plasma.

Keywords: Plasminogen, plasmin, plasma, fractionation.

Introduction

Plasminogen is one of the compounds derived from human plasma by fractionation. When plasminogen is activated to plasmin, it has the property of lysing fibrinogen, fibrin, and some other proteins 1. In another words, plasmin converts fibrin into soluble products and also hydrolyses some other proteins 2. Congenital plasminogen deficiency is an illness that results in inflamed growth on the mucous membranes such as the eyelids and the inside of the mouth 3. Propagation of the growths is usually triggered by infections or injury, but they may also occur spontaneously in the absence of known triggers 4. The prevalence of congenital plasminogen deficiency is about 1.6 per one million people. A reduction in functional plasminogen results in less produced plasmin to break down fibrin, leading to a buildup of fibrin 5. The excess fibrin and the resulting inflammation of the tissue result in the inflamed growths characteristic of congenital plasminogen deficiency.

Formerly plasmin was given parentally for the treatment of thrombotic disorders. Plasmin can be used in combination with deoxyribonucleic for the debridement of wounds ^{6,7}. Also plasminogen has been investigated as a thrombolytic agent and has been used in conjunction with other blood products in wound sealant preparations ^{8,9}.

Usually, isolation of plasminogen needs complex methods ^{10,11}. Using chromate-graphic techniques have great impact in separation and purification of human plasma derived compounds ¹². Among different chromatographic techniques for isolation of human plasma derived products, anion exchange chromatography and affinity chromatography are more applied ^{13,14}.

Ito et al. have separated Glu-plasminogen, Lys-plasminogen and plasmin using high-performance affinity chromatography on a micro particulate polyvinyl alcohol gel as the supporting material and p-aminobenzamidine as the specific ligand ¹⁵.

Mousavi et al.

In another work Cole and Mertz have prepared crude bovine plasminogen from bovine serum by 30% saturation with ammonium sulphate and they reached a 28-fold purification of crude bovine plasminogen ¹⁶.

The aim of the present study was the separation of human plasminogen by affinity chromatography using gel lysine Sepharose. The staring material which we used for chromatography was fraction I supernatant which we prepared using human plasma fractionation.

Materials and Methods

In the current work fraction I was obtained by adjustment of parameters such as ethanol concentration, pH, ionic strength, temperature, and protein concentration as described before 17. We used normal human plasma as the starting material. By increasing its concentration of ethanol from 0% to 8% in pH 7.2 at -3°C the centrifugation was carried out and the supernatant I was separated from precipitate I. Preparation of plasminogen with fraction I supernatant as raw material was followed by the affinity chromatography technique. First the gel of Heparin-Sepharose was suspended in the buffer and applied to the column. Then the column was washed by the solution of sodium chloride (1 M) to remove all proteins bound to the gel due to ion exchange effects. After removal of all impurities the elution of plasminogen was carried out by applying the solution of ε-aminocaproic acid (0.1 M).

Results

In normal human plasma the concentration of plasminogen is about 200mg/l. In our study we could obtain plasminogen with the concentration of 0.5%., which in comparison with the concentration of plasminogen in normal plasma shows about 25 folds higher concentration.

Figure 1 shows the SDS-PAGE electrophoresis of plasminogen with different concentrations at ¼, ½, and non diluted concentration. The SDS-PAGE electrophoresis showed a strong band at 92 kDa related to plasminogen, which normally has a molecular weight of 90 to 92 kDa. The strongest band marked with 4 is related to non-diluted plasminogen and weaker bands marked with 3 and 2 are related to ½ and ¼ diluted plasminogen respectively.




Figure 1: SDS-PAGE electrophoresis of plasminogen with different concentrations.

Discussion

Blood macromolecules usually can be separated from human plasma by plasma fractionation and in combination with chromatographic method ^{18,19-23}. Plasminogen has been prepared using different methods ²⁴⁻²⁶. Benfeldt et al. studied a procedure for the isolation of plasminogen and plasmin from bovine milk. Pure undegraded plasminogen with N-terminal amino acid sequence and mobility in SDS-page similar to plasminogen isolated from bovine blood was obtained ²⁷.

In another work Cole and Mertz prepared crude bovine plasminogen from bovine serum by 30% saturation with ammonium sulphate, and it was partially purified by isoelectric precipitation at pH^{5.3}, and then it was purified further by chromatography on a column of diethylaminoethyl-cellulose ¹⁶. Additional purification of the column product was obtained by precipitation of plasminogen from solution with potassium phosphate and they reached 28-folds purification of crude bovine plasminogen ¹⁶.

Ito et al. separated Glu-plasminogen, Lys-plasminogen and plasmin by high-performance affinity chromatography on a micro particulate polyvinyl alcohol gel as the supporting material and p-aminobenzamidine as the specific ligand ¹⁵.

In the present study, plasminogen with a concentration of 0.5% was achieved which in comparison with the concentration of plasminogen in normal plasma (200 mg/L) shows about 25-fold higher concentration. Finding of our study was in line with Cole and Mertz who purified bovine plasminogen 28-folds from crude plasminogen ¹⁶.

Conclusion

Using affinity chromatography method a 25-fold increase in plasminogen concentration was achieved which shows this method is a promising method for the preparation of plasminogen from human plasma.

References

- Law RH, Abu-Ssaydeh D, Whisstock JC. New insight into the structure and function of the plasminogen/plasmin system. Curr Opin Struct Biol. 2013;23(6):836-41.
- Castellino FJ, Ploplis VA. Structure and function of plasminogen/plasmin system. Thromb Haemost. 2005; 93(4):647-54.
- 3. Mehta R, Shapiro AD. Plasminoge deficiency. Haemophilia. 2008;14(6):1261-8.
- Cohen J, Cohen S, Cymberknon MC, Gross M, Hirshoren N, Shoseyov D. Laryngeal obstruction in congenital plasminogen deficiency. Pediatr Pulmonol. 2012;47(9):923-5.
- 5. Xue Y, Bodin C, Olsson K. Crystal structure of the native plasminogen reveals an activation-resistant compact conformation. J Thromb Heamost. 2012; 10(7):1385-96.
- Li WY, Chomg SS, Huang EY, Tuan TL. Plasminogen activator/plasmin system: a major plater in wound healing? 2003; 11(4):239-47.
- Roth D, Piekarek M, Paulsson M, Christ H, Bloch W, Krieg T, et al. Plasmin modulates vascular endothelial growth factor-A-mediated angiogenesis during wound repair. Am J Pathol. 2006;168(2):670-84.
- Ebrahimian TG, Squiban C, Roque T, Lugo-Martinez H, Hneino M, Buard V, et al. Plasminogen activator inhibitor-1 controlsw bone marrow-derived cells therapeutic effect through MMP9 signalling: role in physiological and pathological wound healing. Stem

- Cells. 2012; 30(7):1436-46.
- Iwaki T, Tanaka A, Miyawaki Y, Suzuki A, Kobayashi T, Takamatsu J, et al. Life-threatening hemorrhage and prolonged wound healing are remarkable phenotypes manifested by complete plasminogen activator inhibitor-1 deficiency in humans. J Thromb Haemost. 2011; 9(6):1200-6.
- Bian L, Ji X, Hu W. Isolation and purification of recombinant human plasminogen Kringle 5 by liquid chromatography and ammonium sulfate saltingout. Biomed Chromatogr. Biomed Chromatogr. 2014; 28(7):957-65.
- Sgier D, Zuberbuehler K, Pfaffen S, Neri D. Isolation and characterization of an inhibitory human monoclonal antibody specific to the urokinase-type plasminogen activator, uPA. Protein Eng Des Sel. 2010; 23(4):261-9.
- 12. Denizli A. Plasma fractionation: conventional and chromatographic methods for albumin purification. J. Biol. & Chem. 2011; 39(4):315-41.
- 13. Vargas M, Segura A, Herrera M, Villalta M, Angulo Y, Gutierrez JM, et al. Purification of IgG and albumin from human plasma by aqueous two phase system fractionation. Biotechnol Prog. 2012; 28(4):1005-11.
- 14. Lihme A, Hansen MB, Andersen IV, Burnouf T. A novel core fractionation process of human plasma by expanded bed adsorption chromatography. Analytical Biochemistry. 2010; 399(1):102-9.
- 15. Ito N, Noguchi K, Kazama M, Shimura K, Kasai K. Separation of human Glu-plasminogen, Lysplasminogen and plasmin by high-performance affinity chromatography on Asahipak GS gel coupled with p-aminobenzamidine. J Chromatogr. 1985; 348(1):199-204.
- Cole ER, Mertz ET. Studies on plasminogen: I. Further purification of bovine plasminogen. Canadian Journal of Biochemistry and Physiology. 1961;39(9):1419-31.
- 17. Burnouf T. Modern plasma fractionation. Transfusion Medicine Reviews. 2007; 21(2):101-7.
- 18. Wallen P, Wiman B. Characterization of human plasminogen II. Separation and partial characterization of different molecular forms of human plasminogen. Biochemica et Biophysica Acta. 1972; 257(1):122-34.
- 19. Mousavi Hosseini K, Pourmokhtar M, Dinarvand R, Rezvan H, Jalili MA. Preparation of enriched immunoglobulin M and immunoglobulin A from human plasma. Medical Journal of the Islamic Republic of Iran. 2004; 17(4):315-8.

Mousavi et al.

- 20. Yari F, Mousavi Hosseini K. Simultaneous purification and polymerization method for bovine serum albumin preparation. The Italian Journal of Biochemistry. 2007; 56(2):163-5.
- Sahebghadam Lofi A, Adibi B, Mousavihosseini K, Mahmoodi M. Purification and isolation of α1-proteinase inhibitor and determination of its' physiological properties. Modarres Journal of Medical Sciences. 2005;7(2):71-80.
- 22. Mousavi Hosseini K, Nasiri S, Heidari M. Using ethanol and low temperature in separation of albumin from human plasma. Journal of Zanjan University of Medical Sciences and Health Services. 2013; 21(85):65-75.
- 23. Mousavi S, Rezvan H, Pourfatollah A, Mousavi Hosseini K. Improving purification of coagulation FIX using heparin affinity chromatography and its comparison with ion exchange chromatography. The Scientific Journal of Iranian Blood Transfusion Organization. 2005; 2(4):91-8.
- Moorhouse KG, Ricke CA, Chen AB. Electrophoretic separation of recombinant tissue-type plasminogen activator glycoforms: Validation issues for capillary isoelectric focusing methods. Electrophoresis. 1996; 17:423-30.
- 25. Nulkar MW, Darad R, Subramanian M, Pawse AR. Plasminogen activator: Isolation and purification from lymphosarcoma of ascites bearing mice. J. Biosci. 1991; 16(6):223-33.
- Tissot JD, Schneider P, Hauert J, Ruegg M, Kruithof EKO, Bachmann F. Isolation from human plasma of a plasminogen activator identical to urinary high molecular weight urokinase. J. Clin. Invest. 1982; 70:1320-3.
- 27. Benfeldt C, Larsen LB, Rasmussen JT, Andersen PA, Petersen TE. Isolation and characterization of plasminogen and plasmin from bovine milk. International Dairy Journal. 1995; 5(6):577-92.