



ORIGINAL ARTICLE

Health-Related Quality of Life in β Thalassemia Major Children in North of Iran

A Jafari-Shakib¹, A Davoudi-Kiakalaye¹, AA Pour-Fathollah², R Jafari-Shakib³, Z Mohtasham-Amiri^{4*}

¹Guilan Blood Transfusion Organization, Rasht, Iran

²Iranian Blood Transfusion Organization, Tehran, Iran

³Department of Immunology, Guilan University of Medical Sciences, Rasht, Iran

⁴Department of Community and Preventive Medicine, Guilan University of Medical Sciences, Rasht, Iran

ARTICLE INFO

Article History:

Received: 15.08.2016

Accepted: 3.10.2016

Keywords:

β thalassemia major
Health-related quality of life
Physical functioning
Social functioning
Emotional functioning

*Corresponding author:

Z Mohtasham-Amiri, MD, MPH
Professor in Preventive and Social
Medicine,
Fellowship in Geriatrics from Sydney
University,
Medical Faculty of Guilan University,
Rasht, Iran
Tel: +98 131 33368773
Fax: +98 131 33753836
Email: mohtashamaz@yahoo.com
mohtasham@gums.ac.ir

ABSTRACT

Background: Advances in treatment of thalassemia major have improved the life expectancy of the patients and therefore their quality of life as other chronic diseases. This study was conducted to assess health-related quality of life in these patients in Guilan province.

Methods: In a cross-sectional study, thirty-one children, aged 8-12 years, with β -thalassemia major were interviewed in Guilan, northern Iran, from January to March 2016. Source of data were health centers of the province and its satellite centers, blood transfusion organizations, general hospitals and private clinics. Health-related quality of life was assessed by using PedsQL™ questionnaire. The Questionnaire was completed at baseline by all patients and their parents. T and Chi-square tests were used as appropriated.

Results: Of the 31 children, 58.1% were girls. Total summary score in children was 75.9 ± 20.1 . Physical, Emotional, social, school and psychosocial functioning scores were 70.6 ± 24 , 73.3 ± 22.9 , 85.9 ± 21 , 74.1 ± 21.5 , 77.7 ± 19.7 , respectively. None of the children underwent splenectomy. Sex, Serum ferritin and hemoglobin levels did not show any association with quality of life in this study.

Conclusion: Although quality of life in these patients was acceptable, HRQOL showed lower scores in comparison to the healthy population. It seems more social and familial support for increasing the quality of life of these children is surely needed.

Please cite this article as: Jafari-Shakib A, Davoudi-Kiakalaye A, Pour-Fathollah AA, Jafari-Shakib R, Mohtasham-Amiri Z. Health-Related Quality of Life in β Thalassemia Major Children in North of Iran. IJBC 2016; 8(4): 108-111.

Introduction

Thalassemia is the most common hereditary anemia in the world with approximately 300 million carriers.¹ This hemoglobinopathy is more prevalent in East Mediterranean region, Southeast Asia and India.²⁻⁵ Thalassemia also is a major health problem in the northern and southern provinces of Iran. Guilan is one of the most prevalent areas of this disease with about 10% of whole thalassemia population.⁶

Although thalassemia major is fatal if not treated

appropriately, due to advances in diagnostic methods and treatment measures during the past decade, prognosis and survival rate of these patients has improved significantly.⁷⁻¹⁰

One of the concerns by improving survival of these people as other chronic diseases is their health-related quality of life. Thalassemia is a chronic disease that presents a diverse range of serious clinical and psychological challenges. These challenges not only do affect on the patients' physical, emotional and social functioning but

also their quality of life and relationship with their family.⁶ For chronic diseases such as thalassemia, where a cure is not attainable and lifelong treatment is needed, HRQOL is likely to be an essential outcome and the allocation of health care resources seem mandatory.

Numerous studies in different geographical areas on quality of life of this group of patients have shown variable results depending on their living environment, access to medical services, available social and familial supports.^{6,11-18}

Thalassemia screening program begun in 1997 in Iran as other countries in the middle-east region resulting in reduction of birth rate of β thalassemia major patients. On the other hand, improving access to health care systems and new treatment modalities have increased life expectancy of these patients. Unfortunately, there is paucity of studies on the quality of life of children suffering from thalassemia major in Iran.¹⁹⁻²⁰ The objective of this study was to determine the actual situation of health-related quality of life in children with β -thalassemia major in Guilan province.

Materials and Methods

In a cross-sectional study, the following organizations were assessed for access to the data of total number of thalassemic patients in Guilan province. All records of non-communicable disease centers, health centers, hospitals and clinics of hematology were extracted and were merged into a single file, using Excel 2012. A total of 45 patients 8-12 years were found of whom thirty-one were interviewed by trained medical team after obtaining written consent from their parents. Socio-demographic data and history of their disease was gathered from their parents. Last level of ferritin and hemoglobin was taken from their records. PedsQLTM for age 8-12 years was used for assessing quality of life of these children. Persian version of this questionnaire had already been validated.²¹ The internal consistency reliability of the PedsQL 4.0 Generic Core Scale approached 0.90 for self-report.²²

The aim of the study was explained to the children and their parents. Participation in study was optional; therefore, verbal consent was taken from all of them. This questionnaire consisted of 23 questions in Likert scale. The PedsQL 4.0 encompasses the essential core domains for pediatric HRQOL measurement: 1) Physical functioning (8 items), 2) Emotional functioning (5 items), 3) Social functioning (5 items) and 4) School functioning (5 items). In the "PedsQLTM Generic Core Scales", for ease of interpretability, items are reversed scored and linearly transformed to a 0-100 scale; the higher points reflect the

higher quality of life. To create the psychosocial health summary score, the mean is computed as the sum of the items divided by the number of items answered in the emotional, social, and school functioning scales. The "Physical Health Summary Score" is the same as the "Physical Functioning Scale Score". To create the Total Scale Score, the mean is computed as the sum of all the items divided by the items answered over all the scales.²²

Data analysis was done by SPSS software, version 18 using description statistics, T and Chi-square tests.

Results

Forty-five children aged 8-12 years with β -thalassemia major living in Guilan province, northern Iran, were enrolled. Of these, 31 patients participated in this study during study period. 18 children (58.1%) were girls and most (64.5%) lived in rural areas. The most common educational level of the parents was diploma or lower. All the patients were receiving iron chelating medications. None of the patients had undergone splenectomy or bone marrow transplant. Age at diagnosis was from 2-6 months old. Mean household income was 270±109 US dollars (range: 67-467 US dollars). The average number of blood transfusions per year was 12 times. Mean hemoglobin and ferritin levels were 8±0.9 g/dl (5.9-9.7 g/dl with median level of 8.1 g/dl) and 1377.5±990 ng/ml (250-4019 ng/ml with median level of 1100 ng/ml), respectively.

Total score in children was 75.9±20.1. Physical functioning, Emotional functioning, Social functioning, and School functioning and "Psychosocial Health Summary Scores" were 70.6±24, 73.3±22.9, 85.9±21, 74.1±21.5, 77.7±19.7, respectively (table 1). Sex, serum ferritin and hemoglobin levels did not show any association with quality of life of the patients.

Discussion

Thalassemia is the most common hereditary anemia in Iran. Treatment measures include allogeneic hematopoietic transplantation if a suitable donor is available or lifelong regular transfusion, iron chelation, splenectomy in occasional cases and supportive measures in terms of growth and nutrition of the patients.

With major improvements in treatment of thalassemic patients in recent years, survivals of the patients have increased and their quality of life has become a major concern for health policy makers and physicians. Studies in different geographical areas have shown that quality of life of the patients highly depend on socio-economic status, living environment, access to medical services and social and family supports.

Table 1: Mean PedsQL 4.0 Generic Quality of Life Scores in patients

Domain	Mean±SD	Median	Min-Max
Physical Functioning	70.6±24	71.8	0-100
Emotional Functioning	73.3±22.9	80	0-100
Social Functioning	85.9±21	90	0-100
School Functioning	74.1±21.5	75	0-100
Psychosocial Health Summary	77.7±19.7	78.3	0-100
Total summary score	75.9±20.1	77	0-100

Previous study illustrated that children above 6 years of age can tell their own health status.¹² HRQOL in our patients indicated higher scores and better conditions in comparison to studies of Ismail in Malaysia,¹² Cheuk in Hong Kong,²³ and Garaibeh in Jordan.¹⁶ However, the scores achieved in our study were similar to other studies.^{5,11,17,24} Based on our results physical and school functioning had lower scores than emotional and social functioning. Frequent absenteeism and low physical performance may be the causes of negative impact of this disease on children's quality of life. It seems that more educational programs for teachers and students could make them more familiar with this disease and asking teacher to spend more time for these students have an important role in improving their situation. In our study, sex did not have any impact on quality of life, but other studies have shown other results in which women have a lower scale of quality of life.^{25,26}

There are also limited studies which show the influence of household income as a contribution factor in quality of life.^{17,27}; however, our study did not confirm it. Inconsistent with finding of other studies;^{5,28,29} serum hemoglobin levels did not show any association with quality of life of the patients. Appropriate control of the disease in our patients may be the reason for almost acceptable quality of life as more than half of our patients had hemoglobin level of ≥ 8 g/dl. Of course small sample size could be another reason for this inconsistent.

Consistent with a previous study in Thailand⁵ serum ferritin level had not any association with quality of life of the patients although many other studies had shown conflicting results.^{8,30,31} Lower age of patients could explain this controversy that short term iron overload did not cause significant complaints and complications. Meanwhile in this study about 44% of patients had serum ferritin level less than 1000 ng/ml.

It seems that over the economic situation, cultural status and acceptance of disease with other member of family are important factors in quality of life.

Conclusion

This was the first study to provide evidence of HRQOL of children with β -thalassemia major in Iran. We found lower scores compared with healthy population. Our study revealed this population need more attention from Health Ministry, Ministry of Education, school authorities and the society to provide them better quality of life.

Acknowledgement

The authors express their gratitude to Iranian Blood Transfusion Organization for their financial support.

Conflict of Interest: None declared.

References

- Ryan K, Bain BJ, Worthington D, James J, Plews D, Mason A, et al. Significant haemoglobinopathies: guidelines for screening and diagnosis. *Br J Haematol*. 2010; 149(1):35–49. doi: 10.1111/j.1365-2141.2009.08054.x. PubMed PMID: 20067565.
- Fucharoen S, Winichagoon P. Prevention and control of thalassemia in Asia. *Asian Biomed J*. 2010; 1(1). doi: 10.5372/abm.v1i1.91.
- AlHamdan NA, AlMazrou YY, AlSwaidi FM, Choudhry A. Premarital screening for thalassemia and sickle cell disease in Saudi Arabia. *Genet Med*. 2007; 9(6):372–77. doi: 10.1097/GIM.0b013e318065a9e8. PubMed PMID: 17575503.
- Sarper N, Senkal V, Guray F, Sahin O, Bayram J. Premarital hemoglobinopathy screening in Kocaeli, Turkey: a crowded industrial center on the north Coast of Marmara Sea. *Turk J Hematol*. 2009; 26(2):62–6. PubMed PMID: 27265274.
- Thavorncharoensap M, Torcharus K, Nuchprayoon I, Riewpaiboon A, Indaratna K, Ubol BO. Factors affecting health-related quality of life in Thai children with thalassemia. HYPERLINK “https://www.ncbi.nlm.nih.gov/pubmed/20180983” \o “BMC blood disorders.” *BMC Blood Disord*. 2010; 10:1. doi: 10.1186/1471-2326-10-1. PubMed PMID: 20180983. PubMed Central PMCID: PMC2836992.
- Jafroodi M, Davoudi-Kiakalayeh A, Mohtasham-Amiri Z, Pourfathollah AA, Haghbin A. Trend in prevalence of hepatitis C virus infection among β -thalassemia major patients: 10 years of experience in Iran. *Int J Prev Med*. 2015; 6:89. doi: 10.4103/2008-7802.164832. PubMed PMID: 26445636. PubMed Central PMCID: PMC4587076.
- Cao A. Quality of life and survival of patients with beta-thalassemia major. *Haematologica*. 2004; 89(10):1157-9. PubMed PMID: 15477196.
- Borgna-Pignatti C, Rugolotto S, De Stefano P, Zhao H, Cappellini MD, Del Vecchio GC, et al. Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica*. 2004; 89(10):1187-93. PubMed PMID: 15477202.
- Porter J, Davis BA. Monitoring chelation treatment to achieve optimal outcome in the treatment of thalassemia. *Best Pract Res Clin Haematol*. 2000; 15(2):329-68. PubMed PMID: 12401311.
- Zurlo MG, De Stefano P, Borgna-Pignatti C, Di Palma A, Piga A, Melevendi C, et al. Survival and causes of death in thalassaemia major. *Lancet*. 1989; 334(8653):27-30. doi: 10.1016/S0140-6736(89)90264-X. PubMed PMID: 2567801.
- Caocci G, Efficace F, Ciotti F, Roncarolo MG, Vacca A, Piras E, et al. Health related quality of life in Middle Eastern children with beta-thalassemia. HYPERLINK “https://www.ncbi.nlm.nih.gov/pubmed/22726530” \o “BMC blood disorders.” *BMC Blood Disord*. 2012; 12:6. doi: 10.1186/1471-2326-12-6. PubMed PMID: 22726530. PubMed Central PMCID: PMC3496588.
- Ismail A, Campbell MJ, Ibrahim HM, Jones GL. Health related quality of life in Malaysian children with thalassaemia. HYPERLINK “https://www.ncbi.nlm.nih.gov/pubmed/16813662” \o “Health and quality of life outcomes.” *Health Qual Life Outcomes*. 2006; 4:39. doi: 10.1186/1477-7525-4-39.

- PubMed PMID: 16813662. PubMed Central PMCID: PMC1538578. 13.
13. Boonchooduang N, Louthrenoo O, Choeyprasert W, Charoenkwan P. Health-related quality of life in adolescents with thalassemia. *Pediatr Hematol Oncol*. 2015; 32(5):341-8. doi:10.3109/08880018.2015.1033795. PubMed PMID: 26086564.
 14. Saha R, Misra R, Saha I. Health Related Quality of Life and it's Predictors among Bengali Thalassemic Children Admitted to a Tertiary Care Hospital. *Indian J Pediatr*. 2015; 82(10):909-16. doi: 10.1007/s12098-014-1670-6. PubMed PMID: 25712005.
 15. Ammad SA, Mubeen SM, Shah SF, Mansoor S. Parents' opinion of quality of life (QOL) in Pakistani thalassaemic children. *J Pak Med Assoc*. 2011; 61(5):470-3. PubMed PMID: 22204182.
 16. Gharaibeh HF, Gharaibeh MK. Factors influencing health-related quality of life of thalassaemic Jordanian children. *Child Care Health Dev*. 2012; 38(2):211-8. doi: 10.1111/j.1365-2214.2011.01224.x.
 17. Surapolchai P, Satayasai W, Sinlapamongkolkul P, Udomsubpayakul U. Biopsychosocial predictors of health-related quality of life in children with thalassemia in Thammasat University Hospital. *J Med Assoc Thai*. 2010; 93 Suppl 7:S65-75. PubMed PMID: 21298837.
 18. Shaligram D, Girimaji SC, Chaturvedi SK. Psychological problems and quality of life in children with thalassemia. *Indian J Pediatr*. 2007; 74(8):727-30. PubMed PMID: 17785893.
 19. Baghiani moghadam MH, Sharifirad G, Rahaei Z, Baghianimoghadam B, Heshmati H. Health related quality of life in children with thalassaemia assessed on the basis of SF-20 questionnaire in Yazd, Iran: a case-control study. *Cent Eur J Public Health*. 2011; 19(3):165-9. PubMed PMID: 22026294.
 20. Tahmasbi S, Aein F, Karimoei M. B- Thalassemia children and quality of life: Does the present condition justify children's quality of life? *Holistic Nursing and Midwifery Journal*. 2007; 17(1); 23-30. (Persian).
 21. Gheissari A, Farajzadegan Z, Heidary M, Salehi F, Masaeli A, Mazrooei A, et al. Validation of Persian version of PedsQL™ 4.0™ generic core scales in toddlers and children. *Int J Prev Med*. 2012; 3(5):341-50. PubMed PMID: 22701775. PubMed Central PMCID: PMC3374492.
 22. Varni JW. The PedsQL™ scoring algorithm: scoring the pediatric quality of life inventory™ [Internet] 1998-2016. Available from: <http://www.pedsq.org/score.html>
 23. Cheuk DK, Mok AS, Lee AC, Chiang AK, Ha SY, Lau YL, Chan GC. Quality of life in patients with transfusion-dependent thalassemia after hematopoietic SCT. *Bone Marrow Transplant*. 2008; 42(5):319-27. doi: 10.1038/bmt.2008.165. PubMed PMID: 18560410
 24. Clarke SA, Skinner R, Guest J, Darbyshire P, Cooper J, Shah F, et al. Health-related quality of life and financial impact of caring for a child with thalassaemia major in the UK. *Child Care Health Dev*. 2010; 36:118–22. doi: 10.1111/j.1365-2214.2009.01043.x. PubMed PMID: 19961496.
 25. Sobota A, Yamashita R, Xu Y, Trachtenberg F, Kohlbray P, Kleinert DA, et al. Quality of life in thalassemia: a comparison of SF-36 results from the thalassemia longitudinal cohort to reported literature and the US norms. *Am J Hematol*. 2011; 86(1): 92–5. doi:10.1002/ajh.21896. PubMed PMID: 21061309. PubMed Central PMCID: PMC4250926.
 26. Hadi N, Karami D, Montazeri A. health-related quality of life in thalassemia major. *Quarterly Journal Payesh*. 2009; 8(4): 387-93. (Persian).
 27. Haghpanah S, Nasirabadi S, Ghaffarpassand F, Karami R, Mahmoodi M, Parand S, et al. Quality of life among Iranian patients with beta-thalassemia major using the SF-36 questionnaire. *Sao Paulo Med J*. 2013; 131(3):166-72. PubMed PMID: 23903265.
 28. Afsar B, Ozdemir NF, Sezer S, Haberal M. Quality of life is not related with liver disease severity but with anemia, malnutrition, and depression in HCV-infected hemodialysis patients. *Hemodial Int*. 2009; 13(1):62-71. doi: 10.1111/j.1542-4758.2009.00329.x. PubMed PMID: 19210280.
 29. Finkelstein FO, Story K, Firaneck C, Mendelssohn D, Barre P, Takano T, et al. Health-related quality of life and hemoglobin levels in chronic kidney disease patients. *Clin J Am Soc Nephrol*. 2009; 4(1):33-8. doi: 10.2215/CJN.00630208. PubMed PMID: 18987300. PubMed Central PMCID: PMC2615698.
 30. Telfer PT, Prestcott E, Holden S, Walker M, Hoffbrand AV, Wonke B. Hepatic iron concentration combined with long-term monitoring of serum ferritin to predict complications of iron overload in thalassaemia major. *Br J Haematol*. 2000; 110(4):971-7. doi: 10.1046/j.1365-2141.2000.02298.x. PubMed PMID: 11054091.
 31. Cianciulli P. Treatment of iron overload in thalassemia. *Pediatr Endocrinol Rev*. 2008; 6 Suppl 1:208-13. PubMed PMID: 19337180.